arXiv:1007.4786v1 [math-ph] 27 Jul 2010

Effective models for conductance in magnetic fields:

derivation of Harper and Hofstadter models

G. De Nittis* and G. Panati**

* SISSA Scuola Internazionale Superiore di Studi Avanzati, Trieste, Italy
`denittis@sissa.it`

** Dipartimento di Matematica, Università di Roma “La Sapienza”, Roma, Italy
`panati@mat.uniroma1.it`

27th July 2010

Abstract

Some relevant transport properties of solids do not depend only on the spectrum of the electronic Hamiltonian, but on finer properties preserved only by unitary equivalence, the most striking example being the conductance. When interested in such properties, and aiming to a simpler model, it is mandatory to check that the simpler effective Hamiltonian is approximately unitarily equivalent to the original one, in the appropriate asymptotic regime. In this paper, we consider the Hamiltonian of an electron in a 2-dimensional periodic potential (*e.g.* generated by the ionic cores of a crystalline solid) under the influence of a uniform transverse magnetic field. We prove that such Hamiltonian is approximately unitarily equivalent to a Hofstadter-like (resp. Harper-like) Hamiltonian, in the limit of weak (resp. strong) magnetic field. The result concerning the case of weak magnetic field holds true in any dimension. Finally, in the limit of strong uniform magnetic field, we show that an additional periodic magnetic potential induces a non-trivial coupling of the Landau bands.

MSC 2010: 81Q15; 81Q20; 81V70 ; 81S05.

Key words: Hofstadter model, Harper model, magnetic Bloch electron.

Contents

1 Introduction **2**

2 Description of the model **7**

3 Space-adiabatic theory for the Hofstadter regime **9**
3.1 Adiabatic parameter for weak magnetic fields . . . 9
3.2 Separation of scales: the Bloch-Floquet transform . . . 10
3.3 The periodic Hamiltonian and the gap condition . . . 12
3.4 τ-equivariant and special τ-equivariant symbol classes . . . 14
3.5 Semiclassics: quantization of equivariant symbols . . . 16
3.6 Main result: effective dynamics for weak magnetic fields . . . 18
3.7 Hofstadter-like Hamiltonians . . . 23

4 Space-adiabatic theory for the Harper regime **25**
4.1 Adiabatic parameter for strong magnetic fields . . . 25
4.2 Separation of scales: the von Neumann unitary . . . 27
4.3 Relevant part of the spectrum: the Landau bands . . . 27
4.4 Symbol class and asymptotic expansion . . . 30
4.5 Semiclassics: the $\mathcal{O}(\delta^4)$-approximated symbol . . . 32
4.6 Main result: effective dynamics for strong magnetic fields . . . 35
4.7 Harper-like Hamiltonians . . . 42
4.8 Coupling of Landau bands in a periodic magnetic potential . . . 43

A Some technical results **45**
A.1 Self-adjointness of H_{BL} and H_{per} . . . 45
A.2 Band spectrum of H_{per} . . . 46
A.3 The Landau Hamiltonian H_{L} . . . 47

B Canonical transform for fast and slow variables **49**
B.1 The transform $\mathcal{W}$ built "by hand" . . . 49
B.2 The integral kernel of $\mathcal{W}$ . . . 51

1 Introduction

Schrödinger operators with periodic potentials and magnetic fields have been fascinating physicist and mathematicians for the last decades. Due to the competition between the crystal length scale and the magnetic length scale, these operators reveals striking features as fractal spectrum [GPPO00], anomalous localization [OGM08], quantization of the transverse conductance [TKNN82, ASS83, BSE94], anomalous thermodynamic phase diagrams [Avr04, OA01].

Due to its relevance for the Quantum Hall Effect (QHE) [Mor88, Gra07], we focus on the two-dimensional *Bloch-Landau Hamiltonian*

$$\mathsf{H}_{\mathrm{BL}} := \frac{1}{2m}\left(-i\hbar\nabla_r - \iota_q \frac{|q|B}{2c} e_\perp \wedge r\right)^2 + \mathsf{V}_\Gamma(r)\,, \tag{1}$$

acting in the Hilbert space $\mathcal{H}_{\mathrm{phy}} = L^2(\mathbb{R}^2, d^2r)$, $r = (r_1, r_2) \in \mathbb{R}^2$. Here c is the speed of light, $h := 2\pi\hbar$ is the Planck constant, m is the mass and q the charge (positive if $\iota_q = 1$ or negative if $\iota_q = -1$) of the charge carrier, B is the strength of the external uniform time-independent magnetic field, $e_\perp = (0,0,1)$ is a unit vector orthogonal to the sample, and V_Γ is a periodic potential describing the interaction of the carrier with the ionic cores of the crystal. For the sake of a simpler notation, in this introduction we assume that the periodicity lattice Γ is simply $\mathbb{Z}^2$.

While extremely interesting, a direct analysis of the fine properties of the operator H_{BL} is a formidable task. Thus the need to study simpler effective models which capture the main features of (1) in suitable physical regimes, as for example in the limit of weak (resp. strong) magnetic field. The relevant dimensionless parameter appearing in the problem is $h_B := {}^{\Phi_0}/_{Z\Phi_B} \propto B^{-1}$, where $\Phi_0 = {}^{hc}/_{e}$ is the *magnetic flux quantum*, $\Phi_B = \Omega_\Gamma B$ is the flux of the external magnetic field through the unit cell of the periodicity lattice Γ (whose area is Ω_Γ) and $Z = {}^{|q|}/_{e}$ is the *magnitude* of the charge q of the carrier in units of e (the *positron* charge). It is also useful to introduce the reduced constant $\hbar_B := {}^{h_B}/_{2\pi}$.

In the limit of weak magnetic field, $\hbar_B \to \infty$, one expects that the relevant features are captured by the well-known *Peierls' substitution* [Pei33, Har55, Hof76], thus yielding to consider, for each Bloch band $\mathcal{E}_* = \mathcal{E}_*(k_1, k_2)$ of interest, the following effective model: in the Hilbert space $L^2(\mathbb{T}^2, d^2k)$, k being the Bloch momentum and $\mathbb{T}^2 \simeq [0, 2\pi) \times [0, 2\pi)$, one considers the Hamiltonian operator

$$H_{\mathrm{Hof}}\,\psi \;= \mathcal{E}_*\left(k - \left(\frac{\iota_q}{\hbar_B}\right)\frac{1}{2}\, e_\perp \wedge i\nabla_k\right)\,\psi, \qquad\qquad \psi \in L^2(\mathbb{T}^2, dk). \tag{2}$$

In physicists' words, the above Hamiltonian corresponds to replace the variables k_1 and k_2 in $\mathcal{E}_n$ with the symmetric operators (*magnetic momenta*)

$$\mathscr{K}_1 := k_1 + \frac{i}{2}\left(\frac{\iota_q}{\hbar_B}\right)\frac{\partial}{\partial k_2}, \qquad\qquad \mathscr{K}_2 := k_2 - \frac{i}{2}\left(\frac{\iota_q}{\hbar_B}\right)\frac{\partial}{\partial k_1}. \tag{3}$$

Since $[\mathscr{K}_1, \mathscr{K}_2] \neq 0$ the latter prescription is formal and (2) must be defined by an appropriate variant of the Weyl quantization. The rigorous justification of the Peierls' substitution and the definition and the derivation of the Hamiltonian (2) are the content of Section 3.

The use of the operator (2) traces back to the pioneering works of R. Peierls [Pei33] and P. G. Harper [Har55], and its spectrum was extensively studied by D. Hofstadter in the celebrated paper [Hof76], where he specialized to the case $\mathcal{E}_*(k_1, k_2) = 2\cos k_1 + 2\cos k_2$. In view of that, we call **Hofstadter-like Hamiltonian** any operator in the form (2), while the name Hofstadter Hamiltonian is used only for the special case above. However, this nomenclature is far to be unique. For instance M. A. Shubin in [Shu94] names *discrete magnetic Laplacian* the same operator (up to a Fourier transform).

As for the case of a strong magnetic field, $\hbar_B \to 0$, the periodic potential can be considered a small perturbation of the Landau Hamiltonian, which provides the leading order approximation of H_{BL}. To the next order of accuracy in $\hbar_B$, to each Landau level there corresponds an effective Hamiltonian, acting in $L^2(\mathbb{R}, dx)$, given (up to a suitable rescaling of the energy scale) by

$$H_{\mathrm{Har}}\ \psi = \mathsf{V}_\Gamma \left(-i(\iota_q \hbar_B) \frac{\partial}{\partial x}, x \right) \psi, \qquad\qquad \psi \in L^2(\mathbb{R}, dx), \tag{4}$$

where the r.h.s refers to the ordinary $(\iota_q \hbar_B)$-Weyl quantization of the $\mathbb{Z}^2$-periodic function $\mathsf{V}_\Gamma : \mathbb{R}^2 \to \mathbb{R}$. We refer to Section 4 for the definition of the effective Hamiltonian (4). We call **Harper-like Hamiltonian** any operator of the form (4), using the name Harper Hamiltonian for the special case $\mathsf{V}_\Gamma(p, x) = 2\cos(2\pi p) + 2\cos(2\pi x)$.

REMARK 1.1 (nomenclature and historical overview)**.** In a remarkable series of papers [HS88, HS90, HS89a] B. Helffer and J. Sjöstrand studied the relation between the spectrum of the operator (4) and the spectra of a one-parameter family of one-dimensional operators on $\ell^2(\mathbb{Z})$ defined by

$$(h_\beta u)_n := u_{n-1} + u_{n+1} + 2\cos(2\pi\theta n + \beta) u_n, \tag{5}$$

where $\theta \in \mathbb{R}$ is a fixed number (*deformation parameter*) and $\beta \in [0, 2\pi)$ is the parameter of the family. In the work of the French authors the operator defined by (5) is called *Harper operator* (and indeed it was introduced by Harper in [Har55]). However, in the last three decades, the operator (5) has been extensively studied by many authors (see [Las05, Las94] for an updated review) with the name of *almost-Mathieu operator*. To avoid confusion and make the nomenclature clear, we chose to adhere to the most recent convention, using the name *almost-Mathieu operator* for (5). We thus decided to give credits to Harper's work by associating his name to the operator (4). ♦◊

The regime of strong magnetic field was originally investigated by A. Rauh [Rau74, Rau75]. However the correct effective model, the operator (4), was derived firstly by M. Wilkinson [Wil87] and then, rigorously, by J. Bellissard in an algebraic context [Bel88] and by B. Helffer and J. Sjöstrand in [HS89b], inspired by the latter paper. In both these papers the case of weak magnetic field is also considered. In particular, in [HS89b] it is proven that H_{Har} (resp. H_{Hof}) has, locally on the energy axis, the same spectrum and the same density of states of H_{BL}, as $\hbar_B \to 0$ (resp. $\hbar_B \to \infty$).

Beyond the spectrum and the density of states, there are other mathematical properties of H_{BL} which reveal interesting physics, as for example the orbital magnetization [GA03, TCVR05, CTVR06] or the transverse (Hall) conductance (1). These properties are not invariant under a loose equivalence relation as isospectrality, then it is important to show that H_{Har} (resp. H_{Hof}) is approximately *unitarily equivalent* to H_{BL} in the appropriate limit. This is the main goal of this paper. The problem is not purely academic, since there exist examples of two isospectral Schrödinger operators which are however *not* unitarily equivalent and which exhibit different values of the transverse conductance. In particular, in [DFP10b] we prove that the Hofstadter operator and the Harper operator are isospectral but not unitarily equivalent. One concludes that, in the study of the conductance, it is not enough to prove that the effective model is isospectral to the original Hamiltonian.

---

(1) Some authors refer to the "Hall conductivity" rather than to the "Hall conductance". Indeed, in two space dimensions the conductivity (microscopic quantity) coincides exactly with the conductance (macroscopic quantity), so the two concepts are synonymous. In this sense the quantization of the Hall conductivity is a macroscopic quantum phenomenon.

We thus introduce the stronger notion of **unitarily effective model**, referring to the concept of almost-invariant subspace introduced by G. Nenciu [Nen02] and to the related notion of effective Hamiltonian [PST03b, PST03a, Teu03], which we shortly review. Let us focus on the regime of weak (resp. strong) magnetic field and define $\varepsilon = \varepsilon(B) = {}^{1}/_{\hbar_B}$ (resp. $\varepsilon(B) = \hbar_B$) so that $\varepsilon \to 0$ in the relevant limit. Let Π_ε be an orthogonal projector in $\mathcal{H}_{\text{phy}}$ such that, for any $N \in \mathbb{N}, N \leq N_0$ there exist a constant C_N such that

$$\|[\mathbf{H}_{\text{BL}};\, \Pi_\varepsilon]\| \leq C_N\, \varepsilon^N \tag{6}$$

for ε sufficiently small. Then $\operatorname{Ran}\Pi_\varepsilon$ is called an **almost-invariant subspace** [Nen02, Teu03] at accuracy N_0, since it follows by a Duhammel's argument that

$$\|(1-\Pi_\varepsilon)\, \mathrm{e}^{-is\,\mathbf{H}_{\text{BL}}}\, \Pi_\varepsilon\| \leq C_N\, \varepsilon^N\, |s|$$

for every $s \in \mathbb{R}$, $N \leq N_0$. Granted the existence of such a subspace, we call **(unitarily) effective Hamiltonian** a self-adjoint operator H_{eff} acting on a Hilbert space $\mathcal{H}_{\text{ref}}$, such that there exists a unitary $U_\varepsilon : \operatorname{Ran}\Pi_\varepsilon \to \mathcal{H}_{\text{ref}}$ such that for any $N \in \mathbb{N}, N \leq N_0$, one has

$$\|\left(\Pi_\varepsilon\, \mathbf{H}_{\text{BL}} - U_\varepsilon^{-1} H_{\text{eff}}\, U_\varepsilon\right) \Pi_\varepsilon\| \leq C'_N\, \varepsilon^N. \tag{7}$$

The estimates (6) and (7) imply that

$$\|\left(\mathrm{e}^{-is\,\mathbf{H}_{\text{BL}}} - U_\varepsilon^{-1}\, \mathrm{e}^{-is\,H_{\text{eff}}}\, U_\varepsilon\right) \Pi_\varepsilon\| \leq C''_N\, \varepsilon^N |s|. \tag{8}$$

When the macroscopic time-scale $t = \varepsilon s$ is physically relevant, the estimate above is simply rescaled. The triple $(\mathcal{H}_{\text{ref}}, U_\varepsilon, H_{\text{eff}})$ is, by definition, a unitarily effective model for $\mathbf{H}_{\text{BL}}$. To our purposes, it is important to notice that the asymptotic unitary equivalence in (7) assures that the *topological invariants* related with the spectral projections of $\Pi_\varepsilon\, \mathbf{H}^\varepsilon_{\text{BL}}\, \Pi_\varepsilon$ (K-theory, Chern numbers, ...) are equal to those of H_{eff}, for ε sufficiently small [DP09].

In this paper we prove that in the limit $\hbar_B \to \infty$ the Hofstadter-like Hamiltonian (2) provides a unitarily effective model for $\mathbf{H}_{\text{BL}}$ with accuracy $N_0 = 1$, and we exhibit an iterative algorithm to construct an effective model at any order of accuracy $N_0 \in \mathbb{N}$ (Theorem 3.12). As for the limit $\hbar_B \to \infty$, up to a rescaling of the energy, the non-trivial leading order (accuracy $N_0 = 1$) for the effective Hamiltonian is given by the Haper-like Hamiltonian (4). We also exhibit the effective Hamiltonian with accuracy $N_0 = 2$, i.e. up to errors of order $\mathcal{O}(\varepsilon^2)$ (Theorem 4.7). Moreover, due to the robustness of the adiabatic techniques, we can generalize the simple model described by (1) to include other potentials, see (9), including in particular a periodic vector potential $\mathbf{A}_\Gamma$. This terms produces interesting consequences especially in the Harper regime (see Section 4.8) and it could play a relevant role in the theory of orbital magnetization. This kind of generalization is new with respect to both [Bel88] and [HS89b].

Our proof is based on the observation that both the Hofstadter and the Harper regime are space-adiabatic limits, and can be treated in the framework of *space-adiabatic perturbation theory* (SAPT) [PST03b, PST03a], see also [Teu03]. As for the Hofstadter regime, the proof follows ideas similar to the ones in [PST03a]. Our generalization allows however to consider a constant magnetic field (while in [PST03a] the vector potential is assumed in $C^\infty_{\mathrm{b}}(\mathbb{R}^d)$) and to include a periodic vector potential. Moreover the proof extends the one in [PST03a], in view of the use of the special symbol classes defined in Section 3.4. On the contrary, from the discussion of the Harper regime $\hbar_B \to 0$ some new mathematical problems emerge. Then, although the "philosophy" of the proof of Theorem 4.7 is of SAPT-type, the technical part is new as it will be explained in Section 4. Notice that the regime of weak magnetic field can also be conveniently approached by using the *magnetic Weyl quantization* [MP04, MPR05, IMP07, IMP09], a viewpoint which is investigated in [DL10].

For the sake of completness, we summarize some salient aspects of the SAPT. We refer to specific references (e.g. [Teu03]) for a complete exposition. Let H be the Hamiltonian of a generic physical system which acts on the total (or physical) Hilbert space $\mathcal{H}$. For the SAPT to be applicable, three important ingredients needs: (i) a distinction between *fast* and *slow degrees of freedom* which is mathematically expressed by a unitary decomposition of the physical space $\mathcal{H}$ into a product space $\mathcal{H}_{\mathrm{s}} \otimes \mathcal{H}_{\mathrm{f}}$ (or, more generally, a direct integral), the first factor being the *space of the slow degrees of freedom* and the second the *space of the fast degrees of freedom*; $\mathcal{H}_{\mathrm{s}} \cong L^2(\mathcal{M})$ for suitable measure space $\mathcal{M}$ is also required; (ii) a dimensionless *adiabatic parameter* $\varepsilon \ll 1$ that quantifies the separation of scales between the fast and slow degrees of freedom and which measures how far are the slow degrees of freedom to be "classical" in terms of some process of quantization; (iii) a *relevant part of the spectrum* for the fast dynamics which remains separated from the rest of the spectrum under the perturbation caused by the slow degrees of freedom.

A numerical simulation of the spectrum of the Hofstadter operator, as a function of the parameter $\varepsilon = {}^1\!/\!_{\hbar_B}$, leads to a fascinating picture known as Hofstadter butterfly [Hof76]. Since the spectrum has zero measure as a subset of the square, the physically relevant object is its complement, the resolvent set. It has been pointed out by D. Osadchy and J. Avron [OA01] that the open connected regions of the resolvent set (*islands*) can be associated to different thermodynamic phases (at zero temperature) of a gas of non interacting fermions in a periodic potential, with $\varepsilon \propto B$ and the chemical potential as thermodynamic coordinates. The different phases are labeled by an integer (*topological quantum number*), interpreted as the value of the transverse conductance of the system in units of ${}^{e^2}\!/\!_h$ in the limit of weak magnetic field. The latter integers are conveniently visualized by different colors, thus leading to the *colored Hofstadter butterfly* [OA01, Avr04]. With this language in mind, the main result of this paper can be reformulated by saying that the Hofstadter-like and Harper-like Hamiltonians are "colour-preserving effective models" for the original Bloch-Landau Hamiltonian. Thus they describe, though in a distorted and approximated way, some aspects of the thermodynamics of the original system.

Acknowledgments. It is a pleasure to thank Y. Avron, J. Bellissard, G. Dell'Antonio,

M. Măntoiu, D. Masoero, H. Spohn and S. Teufel for many useful discussions, and F. Faure for stimulating comments on a previous version of the paper. We are grateful to B. Hellfer for suggesting useful references. Financial support by the INdAM-GNFM project *Giovane ricercatore 2009* is gratefully acknowledged.

2 Description of the model

A generalized Bloch-Landau Hamiltonian

The Hamiltonian (1) describes the dynamics of particle with mass m and charge q which interacts with the ionic structure of a two dimensional crystal and with an external orthogonal uniform magnetic field. A more general model is provided by the operator

$$\mathsf{H}_{\mathrm{BL}} := \frac{1}{2m}\left[-i\hbar\nabla_r - \frac{q}{c}\mathsf{A}_\Gamma(r) - \frac{q}{c}\mathsf{A}(r)\right]^2 + \mathsf{V}_\Gamma(r) + q\ \Phi(r) \tag{9}$$

still called *Bloch-Landau Hamiltonian* and, with an abuse of notation, still denoted with the same symbol used in (1). The vector-valued function $\mathsf{A} := (\mathsf{A}_1, \mathsf{A}_2)$ is a *vector potential* corresponding to an (orthogonal) *external magnetic field* $\mathsf{B} = \nabla_r \wedge \mathsf{A} = (\partial_1\mathsf{A}_2 - \partial_2\mathsf{A}_1)\ e_\perp$, Φ is a *scalar potential* corresponding to a (parallel) *external electric field* $\mathsf{E} = -\nabla_r\Phi$ and A_Γ and V_Γ are *internal periodic potentials* which describe the electromagnetic interaction with the ionic cores of the crystal lattice. The external vector potential is assumed to have the following structure

$$\mathsf{A}(r) = \mathsf{A}_0(r) + \mathsf{A}_B(r), \tag{10}$$

where A_0 is a bounded function and A_B describes a uniform orthogonal magnetic field of strenght B, i.e. in the symmetric gauge

$$\mathsf{A}_B(r) = \frac{B}{2}\ e_\perp \wedge r = \left(-\frac{B}{2}r_2, \frac{B}{2}r_1\right), \qquad \nabla_r \wedge \mathsf{A}_B = B\ e_\perp, \qquad \nabla_r \cdot \mathsf{A}_B = 0. \tag{11}$$

The evolution of the system is prescribed by the Schrödinger equation

$$i\hbar\frac{d}{ds}\psi(r,s) = \mathsf{H}_{\mathrm{BL}}\ \psi(r,s), \tag{12}$$

where s corresponds to the *microscopical time-scale.*

Mathematical description of the crystal structure

The periodicity of the crystal is described by a *two dimensional lattice* $\Gamma \subset \mathbb{R}^2$ (namely a discrete subgroup of maximal dimension of the Abelian group $(\mathbb{R}^2, +)$), thus $\Gamma \simeq \mathbb{Z}^2$. Let $\{a, b\} \subset \mathbb{R}^2$ be two generators of Γ, i.e.

$$\Gamma = \{\gamma \in \mathbb{R}^2\ :\ \gamma = n_1 a + n_2 b, \quad n_1, n_2 \in \mathbb{Z}\}.$$

The *fundamental* or *Voronoi cell* of Γ is $M_\Gamma := \{r \in \mathbb{R}^2 \mid r = l_1\ a + l_2\ b, \quad l_1, l_2 \in [0,1]\}$ and its area is given by $\Omega_\Gamma = |a \wedge b|$. We fix the orientation of the lattice in such a way that

$\Omega_\Gamma = (a_1 b_2 - a_2 b_1) > 0$. We say that a function $f_\Gamma : \mathbb{R}^2 \to \mathbb{C}$ is Γ-*periodic* if $f_\Gamma(r+\gamma) = f_\Gamma(r)$ for all $\gamma \in \Gamma$ and all $r \in \mathbb{R}^2$. The electrostatic and magnetostatic crystal potentials V_Γ and A_Γ are assumed to be Γ-periodic according to the previous definition.

An important notion is that of *dual lattice* Γ^* which is the set of the vectors $\gamma^* \in \mathbb{R}^2$ such that $\gamma^* \cdot \gamma \in 2\pi\mathbb{Z}$ for all $\gamma \in \Gamma$. Let $\{a^*, b^*\} \subset \mathbb{R}^2$ be defined by the relations $a^* \cdot a = b^* \cdot b = 1$ and $a^* \cdot b = b^* \cdot a = 0$; these vectors are the generators of the lattice Γ^*, i.e.

$$\Gamma^* = \{\gamma^* \in \mathbb{R}^2 \ : \ \gamma^* = m_1 \ 2\pi a^* + m_2 \ 2\pi b^*, \quad m_1, m_2 \in \mathbb{Z}\}.$$

The *Brillouin zone* $M_{\Gamma^*} := \{k \in \mathbb{R}^2 \mid k = k_1 \ a^* + k_2 \ b^*, \quad k_1, k_2 \in [0, 2\pi]\}$ is the fundamental cell of the dual lattice Γ^*. The explicit expressions for the dual generators $\{a^*, b^*\}$ in terms of the basis $\{a, b\}$ is

$$a^* = \frac{e_\perp \wedge b}{|a \wedge b|} = \frac{1}{\Omega_\Gamma}(b_2, -b_1), \qquad b^* = -\frac{e_\perp \wedge a}{|a \wedge b|} = \frac{1}{\Omega_\Gamma}(-a_2, a_1). \tag{13}$$

It follows from (13) that the surface of the Brillouin zone is $\Omega_{\Gamma^*} = (2\pi)^2 |a^* \wedge b^*| = {}^{(2\pi)^2}/_{\Omega_\Gamma}$.

Given a Γ-periodic function f_Γ, we denote its Fourier decomposition as

$$f_\Gamma(r) = \sum_{\gamma^* \in \Gamma^*} f(\gamma^*) \ \mathrm{e}^{i\gamma^* \cdot r} = \sum_{m_1, m_2 \in \mathbb{Z}} f_{m_1, m_2} \ \mathrm{e}^{i2\pi(m_1 \ a^* + m_2 \ b^*) \cdot r}. \tag{14}$$

A $\mathbb{Z}^2$-*periodic* function $f : \mathbb{R}^2 \to \mathbb{C}$ is a function periodic with respect to an orthonormal lattice, namely such that $f(x_1 + 1, x_2) = f(x_1, x_2 + 1) = f(x_1, x_2)$ for all $x_1, x_2 \in \mathbb{R}$. If one replaces the two real variables by $x_1 := a^* \cdot r$ and $x_2 := b^* \cdot r$ one has that $f_\Gamma(r) := f(a^* \cdot r, b^* \cdot r)$ is Γ-periodic in r. Every Γ-periodic function can be obtained in this way.

Assumptions on the regularity of the potentials and self-adjointness

Let us denote by $C^n_{\mathrm{b}}(\mathbb{R}^2, \mathbb{R})$ the space of real-valued n-times differentiable functions (smooth functions if $n = \infty$) with continuous and bounded derivatives up to order n. Concerning the internal potentials A_Γ and V_Γ we need to assume that:

Assumption ($\mathrm{A_s}$) [internal potentials, strong form]. *The* Γ*-periodic potential* V_Γ *and the two components of the* Γ*-periodic vector potential* A_Γ *are functions of class* $C^\infty_{\mathrm{b}}(\mathbb{R}^2, \mathbb{R})$.

Sometime will be enough to consider a weaker version of this assumption, namely:

Assumption ($\mathrm{A_w}$) [internal potentials, weak form]. *The two components of the* Γ*-periodic vector potential* A_Γ *are in* $C^1_{\mathrm{b}}(\mathbb{R}^2, \mathbb{R})$. *The* Γ*-periodic potential* V_Γ *verifies the condition* $\int_{M_\Gamma} |\mathsf{V}_\Gamma(r)|^2 \ d^2 r < +\infty$.

Assumption ($\mathrm{A_w}$) implies that V_Γ is *uniformly locally* L^2 and this implies also that V_Γ is infinitesimally bounded with respect to $-\Delta_r$ (see Theorem XIII.96 in [RS78]). Concerning the external potentials A and Φ, we need to assume that:

Assumption (B) [external potentials]. *The scalar potential* Φ *is of class* $C^\infty_{\rm b}(\mathbb{R}^2,\mathbb{R})$. *The vector potential* $\mathbf{A}$ *consists of a linear term* $\mathbf{A}_B$ *of the form (11) plus a bounded term* $\mathbf{A}_0$ *which is of class* $C^\infty_{\rm b}(\mathbb{R}^2,\mathbb{R})$.

When the external potentials $\mathbf{A}$ and Φ vanish, the Bloch-Landau Hamiltonian (9) reduces to the *periodic Hamiltonian* (or *Bloch Hamiltonian*)

$$\mathbf{H}_{\rm per} := \frac{1}{2m}\left[-i\hbar\nabla_r - \frac{q}{c}\mathbf{A}_\Gamma(r)\right]^2 + \mathsf{V}_\Gamma(r). \tag{15}$$

The domains of self-adjointness of $\mathbf{H}_{\rm BL}$ and $\mathbf{H}_{\rm per}$ are described in the following proposition. Its proof, together with some basic notion about the *Sobolev space* $\mathcal{H}^2(\mathbb{R}^2)$ and the *magnetic-Sobolev space* $\mathcal{H}^2_{\rm M}(\mathbb{R}^2)$, is postponed to Section A.1.

Proposition 2.1. *Let* Assumptions $(\mathrm{A_w})$ *and* (B) *hold true. Then both* $\mathbf{H}_{\rm BL}$ *and* $\mathbf{H}_{\rm per}$ *are essentially self-adjoint operators on* $L^2(\mathbb{R}^2, d^2r)$ *with common domain of essential self-adjointness the space of smooth functions with compact support* $C^\infty_{\rm c}(\mathbb{R}^2,\mathbb{C})$. *Moreover the domain of self-adjointness of* $\mathbf{H}_{\rm per}$ *is* $\mathcal{H}^2(\mathbb{R}^2)$ *while the domain of self-adjointness of* $\mathbf{H}_{\rm BL}$ *is* $\mathcal{H}^2_{\rm M}(\mathbb{R}^2)$.

3 Space-adiabatic theory for the Hofstadter regime

3.1 Adiabatic parameter for weak magnetic fields

The SAPT for a Bloch electron developed in [PST03a] is based on the existence of a separation between the *microscopic space scale* fixed by the lattice spacing $\ell := \sqrt{\Omega_\Gamma}$, and a *macroscopic space scale* fixed by the scale of variation of the "slowly varying" external potentials. The existence of such a separation of scales is expressed by introducing a dimensionless parameter $\varepsilon \ll 1$ (*adiabatic parameter*) to control the scale of variation of the vector potential and the scalar potential Φ appearing in (9), namely by setting $\mathbf{A} = \mathbf{A}(\varepsilon r)$ and $\Phi = \Phi(\varepsilon r)$. In particular the external magnetic and electric fields are weak compared to the fields generated by the ionic cores.

It is useful to rewrite the (ε-dependent) Hamiltonian (9) in a dimensionless form. The microscopic unit of length being ℓ, we introduce the dimensionless position vector $x := {}^{r}/_{\ell}$ and the dimensionless gradient $\nabla_x = \ell\nabla_r$. Moreover, since the vector potential has the dimension of a length times a magnetic field, then $A(\varepsilon x) := {}^{\varepsilon}/_{\ell B}\,\mathbf{A}(\varepsilon\ell x)$ is a dimensionless function, with B a dimensional constant which fixes the order of magnitude of the magnetic field due to the external vector potential $\mathbf{A}$. Similarly for $\mathbf{A}_\Gamma$ (with $\varepsilon = 1$). Factoring out the dimensional constants one finds

$$H_{\rm BL} := \frac{1}{\mathcal{E}_0}\mathbf{H}_{\rm BL} = \frac{1}{2}\left[-i\nabla_x - \underbrace{\frac{q\Omega_\Gamma B_\Gamma}{c\hbar}}_{=:\hbar_\Gamma^{-1}} A_\Gamma(x) - \iota_q \underbrace{\frac{|q|\Omega_\Gamma B}{c\hbar}}_{=\hbar_B^{-1}} \frac{1}{\varepsilon} A(\varepsilon x)\right]^2 + V_\Gamma(x) + \phi(\varepsilon x), \tag{16}$$

where $\mathcal{E}_0 := \hbar^2/m\Omega_\Gamma$ is the *natural unit of the energy* fixed by the problem, $V_\Gamma(x) := 1/\mathcal{E}_0\, \mathsf{V}_\Gamma(\ell x)$ and $\phi(\varepsilon x) := q/\mathcal{E}_0\ \Phi(\varepsilon \ell x)$ are both dimensionless quantities. The constant $\hbar_\Gamma$ will play no particular role in the rest of this paper, so it is reabsorbed into the definition of the dimensionless vector potential A_Γ, i.e. formally $\hbar_\Gamma = 1$.

Comparing the dimensional Hamiltonian (16) with the original Hamiltonian (9) , or observing that the strenght of the magnetic field goes to zero (at least linearly) with ε, it is physically reasonable to estimate $\varepsilon\hbar_B \propto 1$. This is rigorously true in the case in which the external electromagnetic field is uniform.

The external force due to A and ϕ are of order of ε and therefore have to act over a time of order ε^{-1} to produce a finite change, which defines the macroscopic time-scale. The *macroscopic dimensionless (slow) time-scale* is fixed by $t := \varepsilon\frac{\mathcal{E}_0}{\hbar}s$ where s is the dimensional microscopic *(fast)* time-scale. With this change of scale the Schrödinger equation (12) reads

$$i\varepsilon\frac{d}{dt}\psi = H_{\mathrm{BL}}\ \psi \tag{17}$$

with H_{BL} given by equation (16).

Remark 3.1. Observe that from the definition of the dimensionless periodic potential A_Γ and V_Γ it follows that they are periodic with respect to the transform $x \mapsto x + \gamma/\ell$. This means that A_Γ and V_Γ are periodic with respect to a "normalized" lattice whose fundamental cell has surface 1. ♦◊

3.2 Separation of scales: the Bloch-Floquet transform

To make explicit the presence of the linear term of the external vector potential, we can rewrite the (16) as follows

$$H_{\mathrm{BL}} = \frac{1}{2}\left[-i\nabla - A_\Gamma(x) - A_0(\varepsilon x) - \iota_q\,\frac{1}{2}e_\perp \wedge \varepsilon x\right]^2 + V_\Gamma(x) + \phi(\varepsilon x), \tag{18}$$

where the adiabatic parameter ε expresses the separation between the macroscopic length-scale, defined by the external potentials, and the microscopic length-scale, defined by the internal Γ-periodic potentials. The separation between slow and fast degrees of freedom can be expressed decomposing the physical Hilbert space $\mathcal{H}_{\mathrm{phy}} = L^2(\mathbb{R}^2, d^2x)$ into a product of two Hilbert spaces or, more generally, into a direct integral. To this end, we use the Bloch-Floquet transform [Kuc93]. As in [PST03a] we define the *(modified) Bloch-Floquet transform* $\mathcal{Z}$ of a function $\psi \in \mathcal{S}(\mathbb{R}^2)$ to be

$$(\mathcal{Z}\psi)(k,\theta) := \sum_{\gamma\in\Gamma} \mathrm{e}^{-i(\theta+\gamma)\cdot k}\psi(\theta+\gamma), \qquad (k,\theta) \in \mathbb{R}^2 \times \mathbb{R}^2. \tag{19}$$

Directly from the definition one can check the following periodicity properties:

$$(\mathcal{Z}\psi)(k,\theta+\gamma) = (\mathcal{Z}\psi)(k,\theta) \qquad \forall\ \gamma \in \Gamma \tag{20}$$

$$(\mathcal{Z}\psi)(k+\gamma^*,\theta) = \mathrm{e}^{-i\theta\cdot\gamma^*}(\mathcal{Z}\psi)(k,\theta) \qquad \forall\ \gamma^* \in \Gamma^*. \tag{21}$$

Equation (20) shows that for any fixed $k \in \mathbb{R}^2$, $(\mathcal{Z}\psi)(k,\cdot)$ is a Γ-periodic function and can be seen as an element of $\mathcal{H}_{\mathrm{f}} := L^2(\mathbb{V}, d^2\theta)$ with $\mathbb{V} := \mathbb{R}^2/\Gamma$ a two-dimensional slant torus (*Voronoi torus*). The torus $\mathbb{V}$ coincides with the the fundamental cell M_Γ endowed with the identification of the opposite edges and $d^2\theta$ denotes the (normalized) measure induced on $\mathbb{V}$ by the identification with M_Γ. The Hilbert space $\mathcal{H}_{\mathrm{f}}$ is the *space of fast degrees of freedom*, corresponding to the microscopic scale. Equation (21) involves a unitary representation $\tau : \Gamma^* \longrightarrow \mathscr{U}(\mathcal{H}_{\mathrm{f}})$ of the group of the (dual) lattice translations Γ^* on the Hilbert space $\mathcal{H}_{\mathrm{f}}$. For every $\gamma^* \in \Gamma^*$ the unitary operator $\tau(\gamma^*)$ is the multiplication with $\mathrm{e}^{i\theta\cdot\gamma^*}$. It will be convenient to introduce the Hilbert space

$$\mathcal{H}_\tau := \left\{\psi \in L^2_{\mathrm{loc}}\left(\mathbb{R}^2, d^2\underline{k}, \mathcal{H}_{\mathrm{f}}\right) \ : \ \psi(k-\gamma^*,\cdot) = \tau(\gamma^*)\ \psi(k,\cdot)\right\} \tag{22}$$

equipped with the inner product $\langle\psi;\varphi\rangle_{\mathcal{H}_\tau} := \int_{M_{\Gamma^*}} (\psi(k);\varphi(k))_{\mathcal{H}_{\mathrm{f}}}\ d^2\underline{k}$ where $d^2\underline{k} := \frac{d^2k}{(2\pi)^2}$ is the normalized measure. There is a natural isomorphism from $\mathcal{H}_\tau$ to $L^2\left(M_{\Gamma^*}, d^2\underline{k}, \mathcal{H}_{\mathrm{f}}\right)$ given by restriction from $\mathbb{R}^2$ to M_{Γ^*}, and with inverse given by τ-covariant continuation, as suggested by (21). The Bloch-Floquet transform (19) extends to a unitary map

$$\mathcal{Z} : \mathcal{H}_{\mathrm{phy}} \longrightarrow \mathcal{H}_\tau \simeq L^2\left(M_{\Gamma^*}, d^2\underline{k}, \mathcal{H}_{\mathrm{f}}\right) \simeq L^2(M_{\Gamma^*}, d^2\underline{k}) \otimes \mathcal{H}_{\mathrm{f}}. \tag{23}$$

The Hilbert space $L^2(M_{\Gamma^*}, d^2\underline{k})$ can be seen as the *space of slow degrees of freedom* and in this sense the transform $\mathcal{Z}$ produces a decomposition of the physical Hilbert space according to the existence of fast and slow degrees of freedom.

We need to discuss how differential and multiplication operators behave under $\mathcal{Z}$. Let $Q = (Q_1, Q_2)$ be the multiplication by $x = (x_1, x_2)$ defined on its maximal domain and $P = (P_1, P_2) = -i\nabla_x$ with domain the Sobolev space $\mathcal{H}^1(\mathbb{R}^2)$, then from (19) it follows:

$$\mathcal{Z}\ P\ \mathcal{Z}^{-1} = k \otimes \mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + \mathbb{1}_{L^2(M_{\Gamma^*})} \otimes -i\nabla_\theta, \qquad\qquad \mathcal{Z}\ Q\ \mathcal{Z}^{-1} = i\nabla^\tau_k \tag{24}$$

where $-i\nabla_\theta$ acts on the domain $\mathcal{H}^1(\mathbb{V})$ while the domain of the differential operator $i\nabla^\tau_k$ is the space $\mathcal{H}_\tau \cap \mathcal{H}^1_{\mathrm{loc}}(\mathbb{R}^2, \mathcal{H}_{\mathrm{f}})$, namely it consists of vector-valued distributions which are in $\mathcal{H}^1\left(M_{\Gamma^*}, \mathcal{H}_{\mathrm{f}}\right)$ and satisfy the θ-dependent boundary condition associated with (21). The central feature of the Bloch-Floquet transform is, however, that multiplication operators corresponding to Γ-periodic functions like A_Γ or V_Γ are mapped into multiplication operators corresponding to the same function, i.e.

$$\mathcal{Z}\ A_\Gamma(x)\ \mathcal{Z}^{-1} = \mathbb{1}_{L^2(M_{\Gamma^*})} \otimes A_\Gamma(\theta) \qquad\qquad \mathcal{Z}\ V_\Gamma(x)\ \mathcal{Z}^{-1} = \mathbb{1}_{L^2(M_{\Gamma^*})} \otimes V_\Gamma(\theta). \tag{25}$$

Let $H^{\mathcal{Z}} := \mathcal{Z}\ H_{\mathrm{BL}}\ \mathcal{Z}^{-1}$ be the Bloch-Floquet transform of the Bloch-Landau Hamiltonian (18). According to the relations (24) and (25) one obtains from (18) that

$$H^{\mathcal{Z}} = \frac{1}{2}\left[-i\nabla_\theta + k - A_\Gamma(\theta) - A_0\left(i\varepsilon\nabla^\tau_k\right) - \iota_q \frac{1}{2} e_\perp \wedge \left(i\varepsilon\nabla^\tau_k\right)\right]^2 + V_\Gamma(\theta) + \phi\left(i\varepsilon\nabla^\tau_k\right) \tag{26}$$

with domain of self-adjointness $\mathcal{Z}\mathcal{H}^2_{\mathrm{M}}(\mathbb{R}^2) \subset \mathcal{H}_\tau$, i.e. the image under $\mathcal{Z}$ of the second magnetic-Sobolev space.

3.3 The periodic Hamiltonian and the gap condition

When $\varepsilon = 0$ the Bloch-Landau Hamiltonian (18) reduces to the periodic Hamiltonian

$$H_{\mathrm{per}} = \frac{1}{2}\left[-i\nabla_x - A_\Gamma(x)\right]^2 + V_\Gamma(x)\,. \tag{27}$$

According to (26) the Bloch-Floquet transform maps H_{per} into a fibered operator. In other words, denoting $H^{\mathcal{Z}}_{\mathrm{per}} := \mathcal{Z}\ H_{\mathrm{per}}\ \mathcal{Z}^{-1}$, one has $H^{\mathcal{Z}}_{\mathrm{per}} = \int^{\oplus}_{M_{\Gamma^*}} H_{\mathrm{per}}(k)\ d^2\underline{k}$ where, for each $k \in M_{\Gamma^*}$

$$H_{\mathrm{per}}(k) = \frac{1}{2}\left[-i\nabla_\theta + k - A_\Gamma(\theta)\right]^2 + V_\Gamma(\theta). \tag{28}$$

The operator $H_{\mathrm{per}}(k)$ acts on $\mathcal{H}_{\mathrm{f}} = L^2(\mathbb{V}, d^2\theta)$ with self-adjointness domain $\mathcal{D} := \mathcal{H}^2(\mathbb{V})$ (the second Sobolev space) independent of $k \in M_{\Gamma^*}$. Moreover it is easy to check that the Bloch-Floquet transform induces the following property of periodicity, called *τ-equivariance*:

$$H_{\mathrm{per}}([k] - \gamma^*) = \tau(\gamma^*)\ H_{\mathrm{per}}([k])\ \tau(\gamma^*)^{-1} \in \Gamma^* \qquad\qquad \forall \gamma^* \in \Gamma^*. \tag{29}$$

where the notation $k := [k] - \gamma^*$ denotes the a.e.-unique decomposition of $k \in \mathbb{R}^2$ as a sum of $[k] \in M_{\Gamma^*}$ and $\gamma^* \in \Gamma^*$.

Remark 3.2 (Analiticity)**.** For any $k \in \mathbb{R}^2$, let $I(k)$ be the unitary operator acting on $\mathcal{H}_{\mathrm{f}}$ as the multiplication by $\mathrm{e}^{-i\theta\cdot k}$. Obviously $I(k) = I([k] - \gamma^*) = I([k])\tau(\gamma^*)^{-1}$. A simple computation shows that

$$H_{\mathrm{per}}(k) = I(k)\ H_{\mathrm{per}}(0)\ I(k)^{-1} \tag{30}$$

where the equality holds on the fixed domain of self-adjointness $\mathcal{D} = \mathcal{H}^2(\mathbb{V})$. The τ-equivariance property (29) follows immediately from (30). Moreover from (30) is evident that $H_{\mathrm{per}}(k)$ defines an *analytic family (of type A) in the sense of Kato* (see [RS78] Chapter XII). Finally a short computation shows

$$(\partial_{k_j} H_{\mathrm{per}})(k) = -iI(k)\ [\theta_j; H_{\mathrm{per}}(0)]\ I(k)^{-1} = I(k)\ \left(-i\nabla_\theta - A_\Gamma(\theta)\right)_j\ I(k)^{-1}$$

and $(\partial^2_{k_j} H_{\mathrm{per}})(k) = \mathbb{1}_{\mathcal{D}}$, $(\partial^2_{k_1,k_2} H_{\mathrm{per}})(k) = 0$ on the domain $\mathcal{D}$. ♦◊

The spectrum of H_{per}, which coincides with the spectrum of $H^{\mathcal{Z}}_{\mathrm{per}}$, is given by the union of all the spectra of $H_{\mathrm{per}}(k)$. The following classical results hold true:

Proposition 3.3. *Let V_Γ and A_Γ satisfy* Assumption ($\mathrm{A_w}$)*, then:*

(i) *for all $k \in \mathbb{R}^2$ the operator $H_{\mathrm{per}}(k)$ defined by* (27) *is self-adjoint with domain $\mathcal{D} = \mathcal{H}^2(\mathbb{V})$ and is bounded below;*

(ii) *$H_{\mathrm{per}}(k)$ has compact resolvent and its spectrum is purely discrete with eigenvalues $\mathcal{E}_n(k) \to +\infty$ as $n \to +\infty$;*

(iii) *let the eigenvalues be arranged in increasing order and repeated according to their multiplicity for any $k \in M_{\Gamma^*}$, i.e. $\mathcal{E}_1(k) \leqslant \mathcal{E}_2(k) \leqslant \mathcal{E}_3(k) \leqslant \ldots$ then $\mathcal{E}_n(k)$ is a continuous Γ^*-periodic function of k.*

The above result differs from the standard theory of periodic Schrödinger operators just for the presence of a periodic vector potential A_Γ. Since we were no able to find a suitable reference in the literature, we sketch its proof in Appendix A.1.

We call $\mathcal{E}_n(\cdot)$ the n^{th} *Bloch band* or *energy band.* The corresponding normalized eigenstates $\{\varphi_n(k)\}_{n\in\mathbb{N}} \subset \mathcal{D}$ are called *Bloch functions* and form, for any $k \in M_{\Gamma^*}$, an orthonormal basis of $\mathcal{H}_{\text{f}}$. Notice that, with this choice of the labelling, $\mathcal{E}_n(\cdot)$ and $\varphi_n(\cdot)$ are continuous in k, but generally they are not smooth functions if *eigenvalue crossings* are present.

We say that a family of Bloch bands $\{\mathcal{E}_n(\cdot)\}_{n\in\mathcal{I}}$, with $\mathcal{I} := [I_+, I_-] \cap \mathbb{N}$, is *isolated* if

$$\inf_{k\in M_{\Gamma^*}} \ \text{dist}\left(\bigcup_{n\in\mathcal{I}}\{\mathcal{E}_n(k)\}, \bigcup_{j\notin\mathcal{I}}\{\mathcal{E}_j(k)\}\right) = C_g > 0. \tag{31}$$

The existence of an isolated part of the spectrum is a necessary ingredient for an adiabatic theory. We introduce the following:

Assumption (C) [constant gap condition]. *The spectrum of H_{per} admits a family of Bloch bands $\{\mathcal{E}_n(\cdot)\}_{n\in\mathcal{I}}$ which is isolated in the sense of* (31)*.*

Let $P_\mathcal{I}(k)$ be the spectral projector of $H_{\text{per}}(k)$ corresponding to the family of eigenvalues $\{\mathcal{E}_n(k)\}_{n\in\mathcal{I}}$, then $P_\mathcal{I}^\mathcal{Z} := \int_{M_{\Gamma^*}}^{\oplus} P_\mathcal{I}(k)\ d^2\underline{k}$ is the projector on the isolated family of Bloch bands labeled by $\mathcal{I}$. In terms of Bloch functions (using the Dirac notation), one has that $P_\mathcal{I}(\cdot) = \sum_{n\in\mathcal{I}} |\varphi_n(\cdot)\rangle\langle\varphi_n(\cdot)|$. However, in general, $\varphi_n(\cdot)$ are not smooth functions of k at eigenvalue crossing, while $P_\mathcal{I}(\cdot)$ is a smooth function of k because of the gap condition. Moreover, from the periodicity of $H_{\text{per}}(\cdot)$, one argues $P_\mathcal{I}([k]-\gamma^*) = \tau(\gamma^*)\ P_\mathcal{I}([k])\ \tau(\gamma^*)^{-1}$. In general the smoothness of $P_\mathcal{I}(\cdot)$ is not enough to assure the existence of family of orthonormal basis for the subspaces $\text{Ran}P_\mathcal{I}(\cdot)$ which varies smoothly (or only continuously) with respect to $k \in M_{\Gamma^*}$. Then we need the following assumption.

Assumption (D) [smooth frame]. *Let $\{\mathcal{E}_n(\cdot)\}_{n\in\mathcal{I}}$ be a family of Bloch bands ($|\mathcal{I}| = m > 1$). We assume that there exists an orthonormal basis $\{\psi_n(\cdot)\}_{j=1}^m$ of $\text{Ran}P_\mathcal{I}(\cdot)$ whose elements are smooth and (left) τ-covariant with respect to k, i.e. $\psi_j(\cdot - \gamma^*) = \tau(\gamma^*)\psi_j(\cdot)$ for all $j = 1, \ldots, m$ and $\gamma^* \in \Gamma^*$.*

Note that it is not required that $\psi_j(k)$ is an eigenfunction of $H_{\text{per}}(k)$. However, in the special but important case in which the family of bands consist of a single isolated m-fold degenerate eigenvalue, i.e. $\mathcal{E}_n(k) = \mathcal{E}_*(k)$ for every $n = 1, \ldots, m$, then the Assumption (D) is equivalent to the existence of an orthonormal basis consisting of smooth and τ-covariant Bloch functions.

Remark 3.4 (Time-reversal symmetry breaking)**.** As far as low dimensional models are concerned ($d \leq 3$), Theorem 1 in [Pan07] assures that Assumption (D) is true whenever the Hamiltonian H_{per} is invariant with respect to the *time-reversal symmetry*, which is implemented in the Schrödinger representation by the complex conjugation operator. However, the term $A_\Gamma \neq 0$ in H_{per} generically breaks the time reversal symmetry. Therefore, to consider also the effects due to a periodic vector potential, we need to assume the existence of a smooth family of frames. Anyway is opinion of the authors that the result in [Pan07] can be extended to the case of a periodic vector potential, at least assuming that A_Γ is small in a suitable sense. ♦◊

Let k_0 be a fixed point in M_{Γ^*} and define the projection $\pi_{\mathrm{r}} := P_{\mathcal{I}}(k_0)$. If the Assumption (C) holds true then dim π_{r} = dim $P_{\mathcal{I}}(k)$ for all $k \in \mathbb{R}^2$. Let $\{\chi_n\}_{j=1}^m$ be an orthonormal basis for Ran π_{r} and define a unitary map

$$u_0(k) := \widetilde{u}_0(k) + u_0^{\perp}(k), \qquad \text{with} \qquad \widetilde{u}_0(k) := \sum_{1\leq j\leq \ell} |\chi_j\rangle\langle\psi_j(k)|, \tag{32}$$

which maps Ran $P_{\mathcal{I}}(k)$ in Ran π_{r}. The definition of this unitary is not unique because the freedom in the choice of the frame and of the orthogonal complement $u_0^{\perp}(k)$. From the definition and the τ-covariance of $\psi_j(\cdot)$ one has that $u_0(k)\ P_{\mathcal{I}}(k)\ u_0(k)^{-1} = \pi_{\mathrm{r}}$ and $u_0([k]-\gamma^*) = u_0([k])\ \tau(\gamma^*)^{-1}$ (*right τ-covariance*).

3.4 τ-equivariant and special τ-equivariant symbol classes

Proposition 3.3 shows that for all $k \in \mathbb{R}^2$, the operator $H_{\mathrm{per}}(k)$ defines an unbounded self-adjoint operator on the Hilbert space $\mathcal{H}_{\mathrm{f}}$ with dense domain $\mathcal{D} := \mathcal{H}^2(\mathbb{V})$. However the domain $\mathcal{D}$ can be considered itself as a Hilbert space with respect to the Sobolev norm $\|\cdot\|_{\mathcal{D}} := \|(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \Delta_\theta)\cdot\|_{\mathcal{H}_{\mathrm{f}}}$ and so $H_{\mathrm{per}}(k)$ can be seen as a bounded linear operator from $\mathcal{D}$ to $\mathcal{H}_{\mathrm{f}}$, i.e. as an element of the Banach space $\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})$. The map $\mathbb{R}^2 \ni k \mapsto H_{\mathrm{per}}(k) \in \mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})$ is a special example of a *operator-valued symbol*. For a summary about the theory of the Weyl quantization of vector-valued symbols, we refer to Appendices A and B in [Teu03]. In what follows we will need the following definition.

Definition 3.5 (Hörmander symbol classes)**.** *A* symbol *is any map F from the (cotangent) space $\mathbb{R}^2\times\mathbb{R}^2$ to the Banach space $\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})$, i.e. $\mathbb{R}^2\times\mathbb{R}^2 \ni (k,\eta) \mapsto F(k,\eta) \in \mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})$. A function $w:\mathbb{R}^2\times\mathbb{R}^2 \to [0,+\infty)$ is said to be an* order function *if there exists constants $C_0>0$ and $N_0>0$ such that*

$$w(k,\eta) \leqslant C_0\left(1+|k-k'|^2+|\eta-\eta'|^2\right)^{\frac{N_0}{2}} w(k',\eta') \tag{33}$$

for every $(k,\eta),(k',\eta') \in \mathbb{R}^2\times\mathbb{R}^2$. A symbol $F \in C^\infty(\mathbb{R}^2\times\mathbb{R}^2, \mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$ is an element of the (Hörmander) symbol class *$S^w(\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$ with order function w, if for every $\alpha,\beta\in\mathbb{N}^2$ there exists a constant $C_{\alpha,\beta}>0$ such that $\|(\partial_k^\alpha\partial_\eta^\beta F)(k,\eta)\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} \leqslant C_{\alpha,\beta}\, w(k,\eta)$ for every $(k,\eta)\in\mathbb{R}^2\times\mathbb{R}^2$.*

According to the previous definition, the vector-valued map $H_{\mathrm{per}}(\cdot)$ defines a Hörmander symbol constant in the η-variables and with order function $v(k,\eta) := 1+|k|^2$ (see the proof of Proposition 3.10 below). However, as showed by equation (29), the symbol $H_{\mathrm{per}}(\cdot)$ satisfies an extra condition of periodicity.

Definition 3.6 (τ-equivariant symbols)**.** *Let Γ^* be a two dimensional lattice (the dual lattice defined in Section 2 for our aims) and $\tau:\Gamma^*\to\mathscr{U}(\mathcal{H}_{\mathrm{f}})$ the unitary representation* *defined in Section 3.2. Denote by* $\widetilde{\tau} := \tau|_{\mathcal{D}}$ *the bounded-operator* [(2)] *representation of* Γ^* *in* $\mathcal{D}$. *A symbol* $F \in S^w(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}}))$ *is said to be* τ-equivariant *if*

$$F(k - \gamma^*, \eta) = \tau(\gamma^*)\ F(k, \eta)\ \widetilde{\tau}(\gamma^*)^{-1} \qquad \forall\ \gamma^* \in \Gamma^*, k \in \mathbb{R}^2.$$

The space of τ*-equivariant symbols is denoted as* $S^w_\tau(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}}))$.

For the purposes of this work, it is convenient to focus on special classes of symbols. By considering the *kinetic momentum function* $\mathbb{R}^2 \times \mathbb{R}^2 \ni (k, \eta) \xmapsto{\kappa} \kappa(k, \eta) := k - A(\eta) \in \mathbb{R}^2$, with A fulfilling Assumption (B), one defines the *minimal coupling map* by

$$(k, \eta) \xmapsto{\jmath_\kappa} \jmath_\kappa(k, \eta) := (\kappa(k, \eta), \eta) \in \mathbb{R}^2 \times \mathbb{R}^2. \tag{34}$$

Definition 3.7 (Special τ-equivariant symbols). *Let* w *be an order function, in the sense of (33). We define*

$$S^w_{\kappa;\tau}(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}})) := \{\widetilde{F} = F \circ \jmath_\kappa\ :\ F \in S^w_\tau(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}}))\}.$$

We refer to $S^w_{\kappa;\tau}(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}}))$ as the class of *special* τ*-equivariant symbols*. The following result shows that special symbols can be considered as genuine τ-equivariant symbols with respect to a modified order function. The key ingredient is the linear growth of the kinetic momentum.

Lemma 3.8. *With the above notations* $S^w_{\kappa;\tau}(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}})) \subset S^{w'}_\tau(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}}))$ *where* $w' := w \circ \jmath_\kappa$.

Proof. If $F \in S^w_\tau(\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}}))$ then also $F \circ \jmath_\kappa$ is τ-equivariant, indeed $\kappa(k-\gamma^*, \eta) = \kappa(k, \eta) - \gamma^*$ and $(F \circ \jmath_\kappa)(k - \gamma^*, \eta) = \tau(\gamma^*)\ (F \circ \jmath_\kappa)(k, \eta)\ \widetilde{\tau}(\gamma^*)^{-1}$. Since $\jmath_\kappa$ is a smooth function, then also the composition $F \circ \jmath_\kappa$ is a smooth function. Observing that $(F \circ \jmath_\kappa)(k, \eta) = F(k - A(\eta), \eta)$ it follows that

$$\begin{aligned}
(\partial_{k_j}(F \circ \jmath_\kappa))(k, \eta) &= ((\partial_{k_j} F) \circ \jmath_\kappa)(k, \eta) \\
(\partial_{\eta_j}(F \circ \jmath_\kappa))(k, \eta) &= ((\partial_{\eta_j} F) \circ \jmath_\kappa)(k, \eta) + \sum_{i=1}^{2} (\partial_{\eta_j} \kappa_i)(k, \eta)\ ((\partial_{k_i} F) \circ \jmath_\kappa)(k, \eta)
\end{aligned}$$

where $\partial_{\eta_j} \kappa_i$ are bounded functions because Assumption (B). From the first equation it follows that

$$\|\partial_{k_j}(F \circ \jmath_\kappa)(k, \eta)\|_{\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}})} \leqslant C_{j,0}\ (w \circ \jmath_\kappa)(k, \eta) \qquad j = 1, 2$$

for suitable positive constants $C_{j,0}$. Similarly the second equation implies

$$\|\partial_{\eta_j}(F \circ \jmath_\kappa)(k, \eta)\|_{\mathscr{B}(\mathcal{D}, \mathcal{H}_{\mathrm{f}})} \leqslant [C_{0,j} + K(C_{1,0} + C_{2,0})](w \circ \jmath_\kappa)(k, \eta).$$

where $K > 0$ is a bound for the functions $\partial_{\eta_j} \kappa_i$. By an inductive argument on the number of the derivatives one can proof that the derivatives of $F \circ \jmath_\kappa$ are bounded by the function $w' := w \circ \jmath_\kappa$. To complete the proof we need to show that w' is an order function according to Definition 3.5. This follows by a simple computation using the fact that κ has a linear growth in k and η. ■

[(2)] Clearly $\tau(\gamma^*)$ acts as an invertible bounded operator on the space $\mathcal{D}$, but it is no longer unitary with respect to the Sobolev-norm defined on $\mathcal{D}$.

In view of Lemma 3.8, all the results of Appendix B of [Teu03] hold true for symbols in $S^{w}_{\kappa;\tau}(\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$ and in particular the quantization of a symbol in $S^{w}_{\kappa;\tau}(\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$ preserves the τ-equivariance and that the pointwise product or the Moyal product of two symbols of order w_1 and w_2 produce a symbol of order $w_1 w_2$ (see [Teu03], Propositions B.3 and B.4).

Remark 3.9 (Notation). In what follows we will use the short notation $F(\kappa;\eta) := (F\circ \jmath_{\kappa})(k,\eta)$ to denote the special symbol $F\circ\jmath_{\kappa}\in S^{w}_{\kappa;\tau}(\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$ related to the τ-equivariant symbols $F\in S^{w}_{\tau}(\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$. We emphasize on the use of the semicolon ";" instead the comma "," and of the symbol of the kinetic momentum κ instead the quasi-momentum k.

♦◊

3.5 Semiclassics: quantization of equivariant symbols

As explained in Section 3.2, the Bloch-Floquet transform $\mathcal{Z}$ provides the separation between the fast degrees of freedom, associated to the Hilbert space $\mathcal{H}_{\mathrm{f}} = L^2(\mathbb{V}, d^2\theta)$, and the slow degrees of freedom, associated to the Hilbert $L^2(M_{\Gamma^*}, d^2\underline{k})$. A fruitful point of view is to consider the slow degrees of freedom "classical" with respect to the "quantum" fast degrees of freedom. Mathematically, this is achived by recognizing that the Hamiltonian $H^{\mathcal{Z}}$ defined in (26) is the *Weyl quantization* of an operator-valued "semiclassical" symbol over the classical phase space $\mathbb{R}^2\times\mathbb{R}^2$. As explained rigorously in the Appendices A and B of [Teu03], the quantization procedure maps an operator-valued symbol $F:\mathbb{R}^2\times\mathbb{R}^2\to\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})$ into a linear operator $\mathrm{Op}_{\varepsilon}(F):\mathcal{S}(\mathbb{R}^2,\mathcal{D})\to\mathcal{S}(\mathbb{R}^2,\mathcal{H}_{\mathrm{f}})$, where $\mathcal{S}(\mathbb{R}^2,\mathcal{H})$ denotes the space of $\mathcal{H}$-valued Schwartz functions. The quantization procedure concerns only the slow degrees of freedom and at a formal level can be identified with the prescription

$$k\longmapsto \mathrm{Op}_{\varepsilon}(k) := \text{multiplication by } k\otimes\mathbb{1}_{\mathcal{D}};\qquad \eta\longmapsto\mathrm{Op}_{\varepsilon}(\eta) := i\varepsilon\nabla_k\otimes\mathbb{1}_{\mathcal{D}}. \tag{35}$$

Let us consider the operator-valued symbol $H_0:\mathbb{R}^2\times\mathbb{R}^2\to\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})$ defined by

$$H_0(k,\eta) := \frac{1}{2}\left[-i\nabla_{\theta} + k - A_{\Gamma}(\theta) - A_0(\eta) - \iota_q\frac{1}{2}e_{\perp}\wedge\eta\right]^2 + V_{\Gamma}(\theta) + \phi(\eta). \tag{36}$$

The symbol H_0 does not depend on ε and in view of Proposition 3.3 it defines an unbounded operator on $\mathcal{H}_{\mathrm{f}}$ with domain of self-adjointness $\mathcal{D} = \mathcal{H}^2(\mathbb{V})$ for all choice of $(k,\eta)\in\mathbb{R}^2\times\mathbb{R}^2$. According to the notation of Section 3.4 and comparing (36) with (28) we can write

$$H_0(k,\eta) = H_{\mathrm{per}}(\kappa(k,\eta)) + \phi(\eta) = (H_{\phi}\circ\jmath_{\kappa})(k,\eta). \tag{37}$$

where $H_{\phi}(k,\eta) := H_{\mathrm{per}}(k) + \phi(\eta)$. As suggested by equation (29), H_{ϕ} is a τ-equivariant symbol. Thus the symbol H_0 is τ-equivariant with respect to the kinetic momentum κ. The following result establishes the exact symbol class for H_0.

Proposition 3.10. *If* Assumption $(\mathrm{A_w})$ *and* (B) *hold true then* $H_0\in S^{v}_{\kappa;\tau}(\mathscr{B}(\mathcal{D};\mathcal{H}_{\mathrm{f}}))$ *with order function* $v(k,\eta) := 1 + |k|^2$.

Proof. Using the result of Lemma 3.8, we only need to show that $H_{\phi}\in S^{v}(\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}}))$. The later claim is easy to verify, indeed the derivative in η are bounded functions, the second derivative in k is a constant and the derivatives of higher order in k are zero. Then we need only to check the growth of the first derivative in k. A simple computation shows that

$$\|(\partial_{k_j} H_\phi)(k,\eta)\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} = \|(\partial_{k_j} H_{\mathrm{per}})(k)\ (\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \Delta_\theta)^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}$$

and since $\partial_{k_j} H_{\mathrm{per}}$ is τ-equivariant (see Remark 3.2), then

$$\|(\partial_{k_j} H_\phi)(k,\eta)\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} = \|(\partial_{k_j} H_{\mathrm{per}})([k])\tau(\gamma^*)^{-1}(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \Delta_\theta)^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}.$$

Observing that $\tau(\gamma^*)$ is the multiplication by $\mathrm{e}^{i\theta\cdot\gamma^*}$ in $\mathcal{H}_{\mathrm{f}}$ and by a simple computation that $(\partial_{k_j} H_{\mathrm{per}})([k])\tau(\gamma^*)^{-1} = \tau(\gamma^*)^{-1}[-2\gamma^*_j + (\partial_{k_j} H_{\mathrm{per}})([k])]$ one has

$$\|(\partial_{k_j} H_\phi)(k,\eta)\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} \leqslant C_1|\gamma^*_j| + \|(\partial_{k_j} H_{\mathrm{per}})([k])\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} \leqslant C_1(|k| + C_3) + C_2$$

where $C_1 = 2\|(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \Delta_\theta)^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}$, $C_2 := \max_{k\in M_{\Gamma^*}} \|(\partial_{k_j} H_{\mathrm{per}})([k])\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})}$ and $|\gamma^*_j| \leqslant |\gamma^*| = |k - [k]| \leqslant |k| + C_3$ with $C_3 := \max_{k\in M_{\Gamma^*}} |k|$. The claim follows observing that $1 + |k| \leqslant 2(1 + |k|^2)$. ■



Figure 1: Structure of the spectrum of $H_0(k,\eta)$. The picture shows schematically a "relevant part of the spectrum", consisting of two energy bands $\{E_*, E_{*+1}\}$, with $E_{*+j}(k,\eta) = \mathcal{E}_{*+j}(\kappa(k,\eta)) + \phi(\eta)$. Notice that we assume only a local gap condition, as stated in (38), while in the picture a stronger condition is satisfied: a gap exists when projecting the relevant bands on the vertical axis.

Equation (37) provides information about the dependence on k and η of the spectrum of H_0. The n^{th} eigenvalue $E_n(k,\eta)$ of the operator $H_0(k,\eta)$ is related to the n^{th} eigenvalue $\mathcal{E}_n(k)$ of the periodic Hamiltonian $H_{\mathrm{per}}(k)$ by the relation $E_n(k,\eta) = \mathcal{E}_n(\kappa(k,\eta)) + \phi(\eta)$. The function $E_n : \mathbb{R}^2 \times \mathbb{R}^2 \to \mathbb{R}$ is still Γ^*-periodic in k but only oscillating with bounded variation in η. Assumption (C) for the family of Bloch bands $\{\mathcal{E}_n(\cdot)\}_{n\in\mathcal{I}}$ immediately implies that

$$\inf_{(k,\eta)\in M_{\Gamma^*}\times\mathbb{R}^2} \operatorname{dist}\left(\bigcup_{n\in\mathcal{I}}\{E_n(k,\eta)\}, \bigcup_{j\notin\mathcal{I}}\{E_j(k,\eta)\}\right) = C_g > 0. \tag{38}$$

This is the relevant part of the spectrum of H_0 which we are interested in.

According to the general theory (see Appendices A and B of [Teu03]), one has that

$$\mathrm{Op}_\varepsilon(H_0) = \frac{1}{2}\left[-i\nabla_\theta + k - A_\Gamma(\theta) - A_0\left(i\varepsilon\nabla_k\right) - \iota_q\frac{1}{2}e_\perp \wedge \left(i\varepsilon\nabla_k\right)\right]^2 + V_\Gamma(\theta) + \phi\left(i\varepsilon\nabla_k\right) \tag{39}$$

defines a linear operator from $\mathcal{S}(\mathbb{R}^2,\mathcal{D})$ in $\mathcal{S}(\mathbb{R}^2,\mathcal{H}_\mathrm{f})$ and by duality it extends to a continuous mapping $\mathrm{Op}_\varepsilon(H_0) : \mathcal{S}'(\mathbb{R}^2,\mathcal{D}) \to \mathcal{S}'(\mathbb{R}^2,\mathcal{H}_\mathrm{f})$ (with an abuse of notation we use the same symbol for the extended operator). The τ-equivariance assures that $\mathrm{Op}_\varepsilon(H_0)\varphi([k]-\gamma^*) = \tau(\gamma^*)\mathrm{Op}_\varepsilon(H_0)\varphi([k])$ (see [Teu03] Proposition B.3). Since $\mathrm{Op}_\varepsilon(H_0)$ preserves τ-equivariance it can then be restricted to an operator on the domain $\mathcal{Z}\mathcal{H}^2_\mathrm{M}(\mathbb{R}^2) \subset \mathcal{S}'(\mathbb{R}^2,\mathcal{D})$ which is the domain of self-adjointness of $H^\mathcal{Z}$, according to (26). To conclude that $\mathrm{Op}_\varepsilon(H_0)$, restricted to $\mathcal{Z}\mathcal{H}^2_\mathrm{M}(\mathbb{R}^2)$, agrees with $H^\mathcal{Z}$ it is enough to recall that $i\nabla^\tau_k$ is defined as $i\nabla_k$ restricted to its natural domain $\mathcal{H}^1\left(\mathbb{R}^2,\mathcal{D}\right) \cap \mathcal{H}_\tau$ and to use the spectral calculus. These arguments justify the following:

Proposition 3.11. *The Hamiltonian $H^\mathcal{Z}$, defined by* (26)*, agrees on its domain of definition with the Weyl quantization of the operator-valued symbol H_0 defined by* (36)*.*

With a little abuse of notation, we refer to this result by writing $H^\mathcal{Z} = \mathrm{Op}_\varepsilon(H_0)$.

3.6 Main result: effective dynamics for weak magnetic fields

Let A_ε and B_ε be ε-dependent (possibly unbounded) linear operators in $\mathcal{H}$. We write $A_\varepsilon = B_\varepsilon + \mathcal{O}_0(\varepsilon^\infty)$ if: for any $N\in\mathbb{N}$ there exist a positive constant C_N such that

$$\|A_\varepsilon - B_\varepsilon\|_{\mathcal{B}(\mathcal{H})} \leq C_N\,\varepsilon^N \tag{40}$$

for every $\varepsilon\in[0,\varepsilon_0)$. Notice that, though the operators are unbounded, the difference is required to be a bounded operator.

We refer to Appendices A and B of [Teu03] for the basic terminology concerning pseudodifferential operators, and in particular as for the notions of *principal symbol*, *asymptotic expansion*, *resummation*, *Moyal product*.

Theorem 3.12. *Let* Assumptions ($\mathrm{A_w}$), (B), (C) *and* (D) *be satisfied and let $\{E_n(\cdot)\}_{n\in\mathcal{I}}$ (with $|\mathcal{I}| = m$) be an isolated family of energy bands for H_0 satisfying condition* (38)*. Then:*

1. Almost-invariant subspace: *there exist an orthogonal projection $\Pi_\varepsilon \in \mathscr{B}(\mathcal{H}_\tau)$, with $\Pi_\varepsilon = \mathrm{Op}_\varepsilon(\pi) + \mathcal{O}_0(\varepsilon^\infty)$ and the symbol $\pi(k,\eta) \asymp \sum_{j=0}^\infty \varepsilon^j\ \pi_j(k,\eta)$ having principal part $\pi_0(k,\eta) = P_\mathcal{I}(k - A(\eta))$, so that*

$$[H^\mathcal{Z};\Pi_\varepsilon] = \mathcal{O}_0(\varepsilon^\infty).$$

In particular for any $N \in \mathbb{N}$ there exist a C_N such that

$$\|(1-\Pi_\varepsilon)\ e^{-i\frac{t}{\varepsilon}H^{\mathcal{Z}}}\ \Pi_\varepsilon\| \leq C_N\ \varepsilon^N\ |t| \tag{41}$$

for ε sufficiently small, $t \in \mathbb{R}$.

2. ***Effective dynamics:*** *let $\mathcal{H}_{\mathrm{ref}} = L^2(M_{\Gamma^*}, d^2\underline{k}) \otimes \mathcal{H}_{\mathrm{f}}$, π_{r} as defined above (32) and $\Pi_{\mathrm{r}} = \mathbb{1}_{L^2(M_{\Gamma^*})} \otimes \pi_{\mathrm{r}} \in \mathscr{B}(\mathcal{H}_{\mathrm{ref}})$. Then there exist a unitary operator*

$$U_\varepsilon : \mathcal{H}_\tau \to \mathcal{H}_{\mathrm{ref}}$$

such that

(i) *$U_\varepsilon = \mathrm{Op}_\varepsilon(u) + \mathcal{O}_0(\varepsilon^\infty)$, where the symbol $u \asymp \sum_{j=0}^{\infty} \varepsilon^j u_j$ has principal part u_0 given by (32) with k replaced by $\kappa(k,\eta)$;*

(ii) $\Pi_{\mathrm{r}} = U_\varepsilon\ \Pi_\varepsilon\ {U_\varepsilon}^{-1}$;

(iii) *posing $\mathcal{K} := \Pi_{\mathrm{r}}\mathcal{H}_{\mathrm{ref}}$, one has*

$$U_\varepsilon\, \Pi_\varepsilon\, H^{\mathcal{Z}}\, \Pi_\varepsilon\, U_\varepsilon^{-1} = H^\varepsilon_{\mathrm{eff}} + \mathcal{O}_0(\varepsilon^\infty) \in \mathscr{B}(\mathcal{K})$$

with $H^\varepsilon_{\mathrm{eff}} = \mathrm{Op}_\varepsilon(h)$ and h a resummation of the formal symbol $u \sharp \pi \sharp H_0 \sharp \pi \sharp u^{-1}$ (thus algorithmically computable at any finite order). Moreover,

$$\|(e^{-i\frac{t}{\varepsilon}H^{\mathcal{Z}}} - U_\varepsilon^{-1}\, e^{-i\frac{t}{\varepsilon}H^\varepsilon_{\mathrm{eff}}}\, U_\varepsilon)\, \Pi_\varepsilon\| \leq C'_N\ \ \varepsilon^N\ (\varepsilon + |t|). \tag{42}$$

Remark 3.13. The previous theorem and the following proof generalize straightforwardly to any dimension $d \in \mathbb{N}$. We prefer to state it only in the case $d = 2$ in view of the application to the QHE and of the comparison with the results in Section 3, the latter being valid only for $d = 2$. ♦◊

Proof of Theorem 3.12

Step 1. Almost-invariant subspace

The proof of the existence of the super-adiabatic projection is very close to the proof of Proposition 1 of [PST03a], so we only sketch the strategy and emphasize the main differences with respect to that proof.

First of all, one constructs a formal symbol $\pi \asymp \sum_{j=0}^{\infty} \varepsilon^j\, \pi_j$ (the *Moyal projection*) such that: (i) $\pi \sharp \pi \asymp \pi$; (ii) $\pi^\dagger = \pi$; (iii) $H_0 \sharp \pi \asymp \pi \sharp H_0$ where $\asymp$ denotes the asymptotic equivalence of formal series.

The symbol π is constructed recursively at any order $j \in \mathbb{N}$ starting from π_0 and H_0. One firstly show the uniqueness of π (see Lemma 2.3. in [PST03b]). The uniqueness allows us to construct π locally, i.e. in a neighborhood of some point $z_0 := (k_0, \eta_0) \in \mathbb{R}^2 \times \mathbb{R}^2$. From the continuity of the map $k \mapsto H_{\mathrm{per}}(k)$ and the condition (31) it follows that there exists a neighborhood $\mathcal{U}_{k_0}$ of k_0 such that for every $k \in \mathcal{U}_{k_0}$ the set $\{\mathcal{E}_n(k)\}_{n\in\mathcal{I}}$ can be enclosed by a positively-oriented circle $\Sigma(k_0) \subset \mathbb{C}$ independent of k. Moreover it is possible to choose $\Sigma(k_0)$ in such a way that: it is symmetric with respect to the real axis; $\mathrm{dist}(\Sigma(k_0), \sigma(H_{\mathrm{per}}(k))) \geqslant \frac{1}{4}C_g$ for all $k \in \mathcal{U}_{k_0}$; $\mathrm{Radius}(\Sigma(k_0)) \leqslant C_r$ is bounded by a constant C_r independent of k_0; $\Sigma(k_0 - \gamma^*) = \Sigma(k_0)$ for all $\gamma^* \in \Gamma^*$.

With the notation of Section 3.4 we have $H_0 = H_\phi \circ \jmath_\kappa$ with $H_\phi(k,\eta) := H_{\mathrm{per}}(k) + \phi(\eta)$. Let $\widetilde{\Lambda}(k_0,\eta_0) := \Sigma(k_0) + \phi(\eta_0)$ denote the translation of the circle $\Sigma(k_0)$ by $\phi(\eta_0) \in \mathbb{R}$ and pose $\Lambda := \widetilde{\Lambda} \circ \jmath_\kappa$. From the smoothness of ϕ it follows that there exists a neighborhood $\mathcal{U}_{z_0} \subset \mathbb{R}^2 \times \mathbb{R}^2$ of z_0 such that $\mathrm{dist}(\Lambda(z_0), \sigma(H_0(z))) \geqslant \frac{1}{4}C_g$ for all $z \in \mathcal{U}_{z_0}$. Moreover $\Lambda(z_0)$ is symmetric with respect to the real axis, has radius bounded by C_r and is Γ^*-periodic in the variable $\kappa = k - A(\eta)$ (see Figure 1).

We proceed by using the *Riesz formula*, namely by posing

$$\pi_j(z) := \frac{i}{2\pi} \oint_{\Lambda(z_0)} d\lambda \; R_j(\lambda, z) \qquad \text{on} \quad \mathcal{U}_{z_0}$$

where $R_j(\lambda,\cdot)$ denotes the j-th term in the *Moyal resolvent* $R(\lambda,\cdot) = \sum_{j=0}^{\infty} \varepsilon^j R_j(\lambda,\cdot)$ (also known as the *parametrix*), defined by the request that

$$(H_0(\cdot) - \lambda \mathbb{1}_{\mathcal{D}}) \,\sharp\, R(\lambda,\cdot) = \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}, \qquad R(\lambda,\cdot) \,\sharp\, (H_0(\cdot) - \lambda \mathbb{1}_{\mathcal{D}}) = \mathbb{1}_{\mathcal{D}} \qquad \text{on} \quad \mathcal{U}_{z_0}.$$

Each term R_j is computed by a recursive procedure starting from $R_0(\lambda,\cdot) := (H_0(\cdot) - \lambda\mathbb{1}_{\mathcal{D}})^{-1}$, as illustrated in [GMS91]. Following [PST03a, equations (30) and (31)] one obtains that

$$R_j(\lambda,z) = -R_0(\lambda,z)\; L_j(\lambda,z) \tag{43}$$

where L_j is the $(j-1)$-th order obstruction for R_0 to be the Moyal resolvent, i.e.

$$(H_0(\cdot) - \lambda\mathbb{1}_{\mathcal{D}}) \,\sharp\, \left(\sum_{n=0}^{j-1} \varepsilon^n \; R_j(\lambda,\cdot) \right) = \mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + \varepsilon^j \; L_j(\cdot) + \mathcal{O}(\varepsilon^{j+1}). \tag{44}$$

At the first order $L_1 = -\frac{i}{2}\{H_0, R_0\}_{k,\eta}$, with $\{\cdot,\cdot\}_{k,\eta}$ the Poisson brackets.

The technical (and crucial) part of the proof is to show that

$$\pi_j \in S^v_{\kappa;\tau}\left(\mathscr{B}(\mathcal{H}_{\mathrm{f}}, \mathcal{D})\right) \cap S^1_{\kappa;\tau}\left(\mathscr{B}(\mathcal{H}_{\mathrm{f}})\right)$$

for all $j \in \mathbb{N}$, with $v(k,\eta) := (1 + |k|^2)$. By means of the recursive construction each $R_j(\lambda,\cdot)$ inherits the special τ-equivariance from the principal symbol $R_0(\lambda,\cdot) = ((H_\phi \circ \jmath_\kappa)(\cdot) - \lambda\mathbb{1}_{\mathcal{D}})^{-1}$. The special periodicity in κ of the domain of integration $\Lambda(\cdot)$ which appears in the Riesz formula assures also the special τ-equivariance of each $\pi_j(\cdot)$.

Since $\|(\partial^\alpha_z \pi_j)(z)\|_\flat \leqslant 2\pi C_r \, \sup_{\lambda\in\Lambda(z_0)} \|\partial^\alpha_z(R_j)(\lambda,z)\|_\flat$ ($\flat$ means either $\mathscr{B}(\mathcal{H}_{\mathrm{f}})$ or $\mathscr{B}(\mathcal{H}_{\mathrm{f}};\mathcal{D})$, $\alpha \in \mathbb{N}^4$ is a multiindex and $\partial^\alpha_z := \partial^{\alpha_1}_{k_1}\partial^{\alpha_2}_{k_2}\partial^{\alpha_3}_{\eta_1}\partial^{\alpha_4}_{\eta_2}$), we need only to prove that $R_j(\lambda,\cdot) \in S^v_{\kappa;\tau}\left(\mathscr{B}(\mathcal{H}_{\mathrm{f}},\mathcal{D})\right) \cap S^1_{\kappa;\tau}\left(\mathscr{B}(\mathcal{H}_{\mathrm{f}})\right)$ uniformly in λ. This is the delicate point of the proof.

First of all, from the definition of $\Lambda(z_0)$ it follows that

$$\|R_0(\lambda,z)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = [\mathrm{dist}(\lambda,\sigma(H_0(z)))]^{-1} \leqslant {}^4\!/_{C_g}$$

uniformly in λ. Let $\sigma \in \mathbb{N}^4$, with $|\sigma| = 1$. One observes that $\partial_z^\sigma R_0(\lambda,z) = -R_0(\lambda,z) N_z^\sigma(\lambda,z)$ with $N_z^\alpha(\lambda,z) := \partial_z^\alpha H_0(z)\ R_0(\lambda,z)$. From the relation

$$\partial_z^\sigma N_z^\alpha = N_z^{\alpha+\sigma} - N_z^\alpha N_z^\sigma$$

and an inductive argument, it follows the *chain rule*

$$\partial_z^\alpha R_0 = R_0\ \sum \omega_{\beta_1\ldots\beta_{|\alpha|}} N_z^{\beta_1}\ldots N_z^{\beta_{|\alpha|}}$$

where $\beta_1,\ldots,\beta_{|\alpha|} \in \mathbb{N}^4$, $|\alpha| := \alpha_1 + \ldots + \alpha_4$, $\omega_{\beta_1\ldots\beta_{|\alpha|}} = \pm 1$ is a suitable sign function and the sum runs over all the combinations of multiindices such that $\beta_1 + \ldots + \beta_{|\alpha|} = \alpha$ with the convention $N_z^0 = \mathbb{1}$. The chain rule implies that $R_0 \in S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ provided that

$$\|N_z^\alpha\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = \|\partial_z^\alpha H_0\ R_0\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C_\alpha \qquad \text{uniformly in } \lambda.$$

The latter condition is true since $\|(\partial_z^\alpha H_0)(k,\eta)\ R_0(\lambda,k,\eta)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant (g\circ \jmath_\kappa)(k,\eta)$, for a suitable $g(k,\eta)$, Γ^*-periodic in k and bounded in η; the latter claim can be checked as in Proposition 3.10.

Similarly to prove that $R_0 \in S^v_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_\mathrm{f},\mathcal{D}))$ we need to show that

$$\|R_0\ N_z^\alpha\|_{\mathscr{B}(\mathcal{H}_\mathrm{f},\mathcal{D})} = \|(\mathbb{1}_{\mathcal{H}_\mathrm{f}} - \Delta_\theta)\ R_0\ N_z^\alpha\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C_\alpha\ v'(\cdot)$$

uniformly in λ. Since N_z^α is bounded on $\mathcal{H}_\mathrm{f}$ it is sufficient to show that $\|(\mathbb{1}_{\mathcal{H}_\mathrm{f}}-\Delta_\theta)R_0(\lambda;z)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C'_\alpha\ v'(z)$. Observe that $\|(\mathbb{1}_{\mathcal{H}_\mathrm{f}}-\Delta_\theta)R_0(\lambda,[\kappa]-\gamma^*;\eta)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = \|(\mathbb{1}_{\mathcal{H}_\mathrm{f}}-\Delta_\theta)\tau(\gamma^*)^{-1}R_0(\lambda,[\kappa];\eta)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$. The commutation relation

$$-\Delta_\theta \tau(\gamma^*)^{-1} = \tau(\gamma^*)^{-1}\left(|\gamma^*|^2 + i2\gamma^*\cdot\nabla_\theta - \Delta_\theta\right)$$

and the straightforward bound

$$\|\left(|\gamma^*|^2 + i2\gamma^*\cdot\nabla_\theta - \Delta_\theta\right)(\mathbb{1}_{\mathcal{H}_\mathrm{f}} - \Delta_\theta)^{-1}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C(1+|\gamma^*|^2) \leqslant C'(1+|\kappa(k,\eta)|^2)$$

imply

$$\|(\mathbb{1}_{\mathcal{H}_\mathrm{f}} - \Delta_\theta)R_0(\lambda,z)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C'_\alpha\ v'(z)\|(\mathbb{1}_{\mathcal{H}_\mathrm{f}} - \Delta_\theta)R_0(\lambda,[\kappa];\eta)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$$

with $v' := v\circ\jmath_\kappa$. Finally observe that

$$\|(\mathbb{1}_{\mathcal{H}_\mathrm{f}} - \Delta_\theta)R_0(\lambda,[\kappa];\eta)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant f([\kappa];\eta) \leqslant C''. \tag{45}$$

The first inequality above follows by an expansion on the Fourier basis, for fixed $[\kappa]$ and η; the second follows from the fact that $[\kappa]$ takes values on a compact set and the explicit dependence on η is through the bounded function ϕ. The bound (45) implies that $R_0 \in S^v_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_\mathrm{f},\mathcal{D})) \cap S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ uniformly in λ.

To prove that $R_j \in S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_\mathrm{f}))$, we observe that for any $\alpha \in \mathbb{N}^d$ one has

$$\partial_z^\alpha R_j(\lambda,z) = R_0(\lambda,z)\ M^\alpha_{z;j}(\lambda,z)$$

where $M^{\alpha}_{z;j}$ is a linear combination of terms which are product of N^{β}_z and $\partial^{\delta}_z L_j$ with $|\beta|, |\delta| \leqslant |\alpha|$. Thus it is sufficient to prove that $L_j \in S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ for every $j \in \mathbb{N}$. The latter claim is proved by induction on $j \in \mathbb{N}$. Referring to (43), one has trivially that $L_1 \in S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$. L_{j+1} is a linear combination of products of N^{α}_z (with $0 \leqslant |\alpha| \leqslant j+1$) and $M^{\beta}_{z,i}$ (with $|\beta| + i = j+1$ and $0 \leqslant i \leqslant j$). Then the induction hypothesis on L_i for all $i = 1, \ldots, j$ implies that L_{j+1} is in $S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$.

Finally observing that $\|\partial^{\alpha}_z R_j\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}},\mathcal{D})} \leqslant \|M^{\alpha}_{z;j}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \|R_0\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}},\mathcal{D})}$ and using the fact that $R_0 \in S^v_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}},\mathcal{D}))$ it follows that $R_j \in S^v_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}},\mathcal{D})) \cap S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ uniformly in λ, for all $j \in \mathbb{N}$.

As explained in Section 3.4, we can apply the result of Proposition B.4 in [Teu03] to special τ-equivariant symbols obtaining $H_0 \,\sharp\, \pi \in S^{v^2}_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$. However the τ-equivariance of $H_0 \,\sharp\, \pi$ and its derivatives implies that the norms are bounded in z, hence $H_0 \,\sharp\, \pi \in S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ which implies by adjointness also $\pi \,\sharp\, H_0 \in S^1_{\kappa;\tau}(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$. By construction $[H^{\mathcal{Z}}; \mathrm{Op}_{\varepsilon}(\pi)] = \mathrm{Op}_{\varepsilon}(H_0 \,\sharp\, \pi - \pi \,\sharp\, H_0) = \mathcal{O}_0(\varepsilon^{\infty})$ where the remainder is bounded in the norm of $\mathscr{B}(\mathcal{H}_{\tau})$.

The operator $\mathrm{Op}_{\varepsilon}(\pi)$ is only approximately a projection, since $\mathrm{Op}_{\varepsilon}(\pi)^2 = \mathrm{Op}_{\varepsilon}(\pi \,\sharp\, \pi) = \mathrm{Op}_{\varepsilon}(\pi) + \mathcal{O}_0(\varepsilon^{\infty})$. We obtain the super adiabatic projection Π_{ε} by using the trick in [NS04]. Indeed, one notices that, for ε sufficiently small, the spectrum of $\mathrm{Op}_{\varepsilon}(\pi)$ does not contain e.g. the points $\{1/2\}$ and $\{3/2\}$. Thus, the formula

$$\Pi_{\varepsilon} = \frac{i}{2\pi} \oint_{|z-1|=1/2} (\mathrm{Op}_{\varepsilon}(\pi) - z)^{-1}. \tag{46}$$

yields an orthogonal projector such that $\Pi_{\varepsilon} = \mathrm{Op}_{\varepsilon}(\pi) + \mathcal{O}_0(\varepsilon^{\infty})$.

Finally, equation (41) follows by observing that $[H^{\mathcal{Z}}; \Pi_{\varepsilon}] = \mathcal{O}_0(\varepsilon^{\infty})$ implies

$$[\mathrm{e}^{-i\frac{t}{\varepsilon}H^{\mathcal{Z}}}; \Pi_{\varepsilon}] = \mathcal{O}_0(\varepsilon^{\infty}|t|)$$

as proved in Corollary 3.3 in [Teu03].

Step 2. Construction of the intertwining unitary

The construction of the intertwining unitary follows as in the proof of Proposition 2 of [PST03a]. Firstly one constructs a formal symbol $u \asymp \sum_{j=0}^{\infty} \varepsilon^j u_j$ such that: (i) $u^{\dagger} \,\sharp\, u = u \,\sharp\, u^{\dagger} = \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}$; (ii) $u \,\sharp\, \pi \,\sharp\, u^{\dagger} = \pi_{\mathrm{r}}$.

The existence of such a symbol follows from a recursive procedure starting from u_0 and using the expansion of $\pi \asymp \sum_{j=0}^{\infty} \varepsilon^j \pi_j$ obtained above. However, the symbol u which comes out of this procedure is not unique.

Since u_0 is *right τ-covariant* (see the end of Section 3.3) in κ, then one can prove by induction that the same is also true for all the symbols u_j and hence for the full symbol u. Finally, since $u_0 \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ one deduces by induction also $u_j \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ for all $j \in \mathbb{N}$. The quantization of this symbol is an element of $\mathscr{B}(\mathcal{H}_{\tau}, \mathcal{H}_{\mathrm{ref}})$ satisfying the following properties:

(i) $\mathrm{Op}_\varepsilon(u)\mathrm{Op}_\varepsilon(u)^\dagger = \mathbb{1}_{\mathcal{H}_{\mathrm{ref}}} + \mathcal{O}_0(\varepsilon^\infty)$,

(ii) $\mathrm{Op}_\varepsilon(u)^\dagger\mathrm{Op}_\varepsilon(u) = \mathbb{1}_{\mathcal{H}_\tau} + \mathcal{O}_0(\varepsilon^\infty)$,

(iii) $\mathrm{Op}_\varepsilon(u)\Pi_\varepsilon\mathrm{Op}_\varepsilon(u)^\dagger = \Pi_{\mathrm{r}} + \mathcal{O}_0(\varepsilon^\infty)$.

Nevertheless $\mathrm{Op}_\varepsilon(u)$ can be modified by an $\mathcal{O}_0(\varepsilon^\infty)$ term using the same technique of Lemma 3.3 (Step II) in [PST03b] to obtain the true unitary U_ε.

Step 3. Effective dynamics

The last step of the proof is identical to the corresponding part (Proposition 3) of [PST03a].

3.7 Hofstadter-like Hamiltonians

We now focus on the special case of a single isolated energy band E_*, i.e. $m = 1$, and we comment on the relation between the effective Hamiltonian, the celebrated Peierls substitution and Hofstadter-like Hamiltonians (see Section 1).

In this special case, $\pi_0(\kappa) = |\psi_*(\kappa)\rangle\langle\psi_*(\kappa)|$ and $u_0(\kappa) = |\chi\rangle\langle\psi_*(\kappa)| + u_0^\perp$ where $\psi_*(k)$ is the eigenvector of $H_{\mathrm{per}}(k)$ corresponding to the eigenvalue $\mathcal{E}_*(k)$. Let $h \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ be a resummation of the formal symbol $u \,\sharp\, \pi \,\sharp\, H_0 \,\sharp\, \pi \,\sharp\, u^{-1}$. A straightforward computation yields

$$h_0 = u_0\,\pi_0\,H_0\,\pi_0\,u_0^\dagger = |\chi\rangle\langle\psi_*|\;|\psi_*\rangle\langle\psi_*|\;H_0\;|\psi_*\rangle\langle\psi_*|\;|\psi_*\rangle\langle\chi| = E_*\,\pi_{\mathrm{r}}.$$

Since π_{r} is one-dimensional, h_0 can be regarded as a scalar-valued symbol with explicit expression

$$h_0(k,\eta) = E_*(k,\eta) = \mathcal{E}_*\left(k - A(\eta)\right) + \phi(\eta).$$

By considering the quantization of the latter, the effective one-band Hamiltonian reads

$$\mathrm{Op}_\varepsilon(h_0) = E_*(k, i\varepsilon\nabla_k) = \mathcal{E}_*\left(k - A(i\varepsilon\nabla_k)\right) + \phi(i\varepsilon\nabla_k). \tag{47}$$

The latter formula corresponds to the momentum-space reformulation of the well-known ***Peierls substitution*** [Pei33, AM76].

To illustrate this point, we specialize to the case of a uniform external magnetic field and zero external electric field, setting $\phi = 0$ and $A_0 = 0$ in (36). The leading order contribution to the dynamics in the almost invariant subspace is therefore given by a bounded operator, acting on the reference Hilbert space $L^2(M_{\Gamma^*}, d^2\underline{k})$, defined as the quantization (in the sense of Section 3.5) of the function $\mathcal{E}_* \circ \jmath_\kappa : (k,\eta) \mapsto \mathcal{E}_*(k - A(\eta))$, defined on $\mathbb{T}^d \times \mathbb{R}^d$.

Loosely speaking, the above procedure corresponds to the following "substitution rule": one may think to quantize the smooth function $\mathcal{E}_* : \mathbb{T}^d \to \mathbb{R}$ by formally replacing the variables (k_1, k_2) with the operators $(\mathscr{K}_1', \mathscr{K}_2')$ defined by

$$\mathscr{K}_1' := k_1 + \frac{i}{2}(\iota_q\varepsilon)\frac{\partial}{\partial k_2}, \qquad \mathscr{K}_2' := k_2 - \frac{i}{2}(\iota_q\varepsilon)\frac{\partial}{\partial k_1}, \tag{48}$$

regarded as unbounded operators acting in $L^2(M_{\Gamma^*}, d^2\underline{k})$. To make this procedure rigorous, one can expand $\mathcal{E}_*$ in its Fourier series, i.e. $\mathcal{E}_*(k) = \sum_{n,m\in\mathbb{Z}} c_{n,m}\ \mathrm{e}^{i2\pi(na+mb)\cdot k}$ and define the ***Peierls' quantization*** of $\mathcal{E}_*$ as the operator obtained by the same series expansion with the phases $\mathrm{e}^{i2\pi(na+mb)\cdot k}$ replaced by the unitary operators $\mathrm{e}^{i2\pi(na+mb)\cdot\mathscr{K}'}$ (the series is norm-convergent, in view of the regularity of $\mathcal{E}_*$). This fixes uniquely the prescription for the quantization.

To streamline the notation, one introduces new coordinates $\xi_1 := 2\pi(a\cdot k)$ and $\xi_2 := 2\pi(b\cdot k)$ such that the function $\mathcal{E}'_*$, $\mathcal{E}'_*(\xi_1,\xi_2) := \mathcal{E}_*(k(\xi))$ becomes $(2\pi\mathbb{Z})^2$-periodic. The change of variables induces a unitary map from $L^2({M_{\Gamma^*}}^2, d^2\underline{k})$ to $L^2(\mathbb{T}^2, d^2\xi)$ which intertwines the operators (48) with the operators (recall $\varepsilon = {}^{2\pi}/_{h_B}$)

$$\mathscr{K}_1 := \xi_1 + i\pi\left(\frac{\iota_q}{h_B}\right)\frac{\partial}{\partial\xi_2}, \qquad \mathscr{K}_2 := \xi_2 - i\pi\left(\frac{\iota_q}{h_B}\right)\frac{\partial}{\partial\xi_1}, \tag{49}$$

so that $2\pi(a\cdot\mathscr{K}') \mapsto \mathscr{K}_1$ and $2\pi(b\cdot\mathscr{K}') \mapsto \mathscr{K}_2$.

Let $F : \mathbb{T}^2 \to \mathbb{C}$ be sufficiently regular that its Fourier series $F(\xi_1,\xi_2) = \sum_{n,m\in\mathbb{Z}} f_{n,m}\mathrm{e}^{i(n\xi_1+m\xi_2)}$ is uniformly-convergent. We define the *Peierls quantization* of F as

$$\widehat{F} := \sum_{n,m\in\mathbb{Z}} f_{n,m}\mathrm{e}^{i(n\mathscr{K}_1+m\mathscr{K}_2)}.$$

Let $\mathcal{U}_0 = \mathrm{e}^{i\mathscr{K}_1}$ and $\mathcal{V}_0 = \mathrm{e}^{i\mathscr{K}_2}$ (*Hofstadter unitaries*), acting on $L^2(\mathbb{T}^2, d^2\xi)$ as

$$(\mathcal{U}_0\psi)(\xi_1,\xi_2) = \mathrm{e}^{i\xi_1}\psi\left(\xi_1, \xi_2 - \pi\frac{\iota_q}{h_B}\right), \qquad (\mathcal{V}_0\psi)(\xi_1,\xi_2) = \mathrm{e}^{i\xi_2}\psi\left(\xi_1 + \pi\frac{\iota_q}{h_B}, \xi_2\right). \tag{50}$$

We regard (50) as the definition of the two unitaries, so there is no need to specify the domain of definition of the generators (49). Thus the Peierls quantization of the function F defines a bounded operator on $L^2(\mathbb{T}^2, d^2\xi)$ given, in terms of the Hofstadter unitaries, by

$$\widehat{F}(\mathcal{U}_0,\mathcal{V}_0) = \sum_{n,m=-\infty}^{+\infty} f_{n,m}\,\mathrm{e}^{i\pi nm\left(\frac{\iota_q}{h_B}\right)}\ \mathcal{U}_0^n\ \mathcal{V}_0^m, \tag{51}$$

where the fundamental commutation relation $\mathcal{U}_0\mathcal{V}_0 = \mathrm{e}^{-i2\pi\left(\frac{\iota_q}{h_B}\right)}\ \mathcal{V}_0\mathcal{U}_0$ has been used. Formula (51) defines a *Hofstadter-like Hamiltonian* with *deformation parameter* ${}^1/_{h_B}$. Indeed, the special case $H_{\mathrm{Hof}} = \mathcal{U}_0 + \mathcal{U}_0^{-1} + \mathcal{V}_0 + \mathcal{V}_0^{-1}$ is (up to a unitary equivalence) the celebrated Hofstadter Hamiltonian [Hof76].

Summarizing, we draw the following

Conclusion 3.14. *Under the assumption of Theorem 4.7, for every* $\mathsf{V}_\Gamma \in L^2_{\mathrm{loc}}(\mathbb{R}^2, d^2r)$, *in the Hofstadter regime* ($h_B \to \infty$), *the dynamics generated by the Hamiltonian* $\mathbf{H}_{\mathrm{BL}}$ (1) *in the subspace related to a single isolated Bloch band, is approximated up to an error of order* ${}^1/_{h_B}$ *(and up to a unitary transform) by the dynamics generated on the reference Hilbert space* $L^2(\mathbb{T}^2, d^2\xi)$ *by a Hofstadter-like Hamiltonian, i.e. by a power series in the Hofstadter unitaries* $\mathcal{U}_0$ *and* $\mathcal{V}_0$ *defined by* (50).

4 Space-adiabatic theory for the Harper regime

4.1 Adiabatic parameter for strong magnetic fields

We now consider the case of a strong external magnetic field. Since we are interested in the limit $B \to +\infty$ we set $\mathbf{A}_0 = 0$ and $\Phi = 0$ in the Hamiltonian (9). By exploiting the gauge freedom, we choose

$$\nabla_r \cdot \mathbf{A}_\Gamma = 0, \qquad\qquad \int_{M_\Gamma} \mathbf{A}_\Gamma(r)\; d^2r = 0, \tag{52}$$

this choice being always possible [Sob97]. Let us denote by $Q_r = (Q_{r_1}, Q_{r_2})$ the multiplication operators by r_1 and r_2 and with $P_r = (P_{r_1}, P_{r_2}) = -i\hbar\nabla_r$. Taking into account conditions (52) and $A_0 = 0$, $\phi = 0$, the Hamiltonian (9) is rewritten as

$$\mathbf{H}_{\mathrm{BL}} = \frac{1}{2m}\left[\left(P_{r_1} + \frac{qB}{2c}Q_{r_2}\right)^2 + \left(P_{r_2} - \frac{qB}{2c}Q_{r_1}\right)^2\right] + \widetilde{\mathsf{V}}_\Gamma(Q_r) + \widetilde{\mathsf{W}}(Q_r) \tag{53}$$

where

$$\widetilde{\mathsf{V}}_\Gamma(Q_r) = \mathsf{V}_\Gamma(Q_r) + \frac{q^2}{2mc^2}|\mathbf{A}_\Gamma(Q_r)|^2 \tag{54}$$

$$\widetilde{\mathsf{W}}(Q_r, P_r) = -\frac{q}{mc}(\mathsf{A}_\Gamma)_1(Q_r)\;\left[P_{r_1} + \frac{qB}{2c}Q_{r_2}\right] - \frac{q}{mc}(\mathsf{A}_\Gamma)_2(Q_r)\;\left[P_{r_2} - \frac{qB}{2c}Q_{r_1}\right] \tag{55}$$

with $(\mathsf{A}_\Gamma)_1$ and $(\mathsf{A}_\Gamma)_2$ the Γ-periodic components of the vector potential $\mathbf{A}_\Gamma$. The first of (52) assures that $\widetilde{\mathsf{W}}$ is a symmetric operator.

It is useful to define two new pairs of canonical dimensionless operators:

$$(\text{fast}) \begin{cases} K_1 := -\dfrac{1}{2\delta}\; b^* \cdot Q_r - \iota_q \dfrac{\delta}{\hbar}\; a \cdot P_r \\ K_2 := \dfrac{1}{2\delta}\; a^* \cdot Q_r - \iota_q \dfrac{\delta}{\hbar}\; b \cdot P_r \end{cases} \qquad (\text{slow}) \begin{cases} G_1 := \dfrac{1}{2}\; b^* \cdot Q_r - \iota_q \dfrac{\delta^2}{\hbar}\; a \cdot P_r \\ G_2 := \dfrac{1}{2}\; a^* \cdot Q_r + \iota_q \dfrac{\delta^2}{\hbar}\; b \cdot P_r \end{cases} \tag{56}$$

where $\delta := \sqrt{\hbar_B} = \sqrt{\Phi_0/2\pi Z \Phi_B}$ according to the notation introduced in Section 1. Since $\delta^2 \propto 1/B$, the limit of strong magnetic field corresponds to $\delta \to 0$. We consider δ as the *adiabatic parameter* in the Harper regime. A direct computation shows that

$$[K_1; K_2] = i\iota_q\; \mathbb{1}_{\mathcal{H}_{\mathrm{phy}}}, \qquad [G_1; G_2] = i\iota_q\; \delta^2\; \mathbb{1}_{\mathcal{H}_{\mathrm{phy}}}, \qquad [K_j; G_k] = 0, \qquad j,k = 1,2. \tag{57}$$

These new variables are important for three reasons:

(a) they make evident a separation of scales between the slow degrees of freedom related to the the dynamics induced by periodic potential and the fast degrees of freedom related to the cyclotron motion induced by the external magnetic field. Indeed, for $\mathsf{V}_\Gamma = 0$, the *fast variables* (K_1, K_2) (the *kinetic momenta*) describe the kinetic energy of the cyclotron motion, while the *slow variables* (G_1, G_2) correspond semiclassically to the center of the cyclotron orbit and are conserved quantities.

(b) The new variables are dimensionless. According to the notation used in Section 3.1 let $H_{\mathrm{BL}} := {}^{1}\!/_{\mathcal{E}_0}\mathsf{H}_{\mathrm{BL}}$ be the dimensionless Bloch-Landau Hamiltonian with $\mathcal{E}_0 := {}^{\hbar^2}\!/_{m\Omega_\Gamma}$ the *natural unit of energy.*

(c) The use of the new variables simplifies the expression of the Γ-periodic functions appearing in H_{BL}. Indeed, $a^* \cdot Q_r = G_2 + \delta\ K_2$ and $b^* \cdot Q_r = G_1 - \delta\ K_1$, hence if f_Γ is any Γ-periodic function one has

$$f_\Gamma(Q_r) = f(G_2 + \delta\ K_2, G_1 - \delta\ K_1) \tag{58}$$

where f is the $\mathbb{Z}^2$-periodic function related to f_Γ.

In terms of the new variables (56), the Hamiltonian H_{BL} reads

$$H_{\mathrm{BL}} = \frac{1}{\delta^2}\ \Xi(K_1, K_2) + V\left(G_2 + \delta\ K_2, G_1 - \delta\ K_1\right) + \frac{1}{\delta}\ W(K_1, G_1, K_2, G_2) \tag{59}$$

where

$$\Xi(K_1, K_2) := \frac{1}{2\Omega_\Gamma}\left[|a|^2 {K_2}^2 + |b|^2 {K_1}^2 - a \cdot b\ \{K_1; K_2\}\right] \tag{60}$$

is a quadratic function of the operators K_1 and K_2 ($\{\cdot;\cdot\}$ denotes the anticommutator), V is the $\mathbb{Z}^2$-periodic function related to the Γ-periodic function ${}^{1}\!/_{\mathcal{E}_0}\widetilde{\mathsf{V}}_\Gamma$ and W denotes the function ${}^{1}\!/_{\mathcal{E}_0}\widetilde{W}$ with respect the new canonical pairs, namely

$$W(K_1, G_1, K_2, G_2) = f_1\left(G_2 + \delta\ K_2, G_1 - \delta\ K_1\right) K_1 - f_2\left(G_2 + \delta\ K_2, G_1 - \delta\ K_1\right) K_2 \tag{61}$$

where f_1 and f_2 are the $\mathbb{Z}^2$-periodic dimensionless functions

$$f_1(a^* \cdot r, b^* \cdot r) := 2\pi\frac{Z\Omega_\Gamma}{\Phi_0}(a^* \cdot \mathbf{A}_\Gamma)(r) \qquad \text{and} \qquad f_2(a^* \cdot r, b^* \cdot r) := 2\pi\frac{Z\Omega_\Gamma}{\Phi_0}(b^* \cdot \mathbf{A}_\Gamma)(r).$$

An easy computation shows that the first gauge condition of (52) is equivalent to

$$\frac{\partial f_1}{\partial x_1}(x_1, x_2) + \frac{\partial f_2}{\partial x_2}(x_1, x_2) = 0. \tag{62}$$

Obviously W is a symmetric operator, since $\widetilde{\mathsf{W}}$ is symmetric.

The problem has a natural time-scale which is fixed by the *cyclotron frequency* $\omega_c = \frac{|q|B}{mc}$. With respect to the (fast) *ultramicroscopic time-scale* $\tau := \omega_c s$, equation (12) becomes

$$i\frac{1}{\delta^2}\frac{\partial}{\partial\tau}\psi = H_{\mathrm{BL}}\psi, \qquad\qquad \delta^2 = \frac{\mathcal{E}_0}{\hbar\omega_c}. \tag{63}$$

Thus the physically relevant Hamiltonian is

$$H^\delta_{\mathrm{BL}} := \delta^2\ H_{\mathrm{BL}} = \Xi(K_1, K_2) + \delta\ W(K_1, G_1, K_2, G_2) + \delta^2\ V\left(G_2 + \delta\ K_2, G_1 - \delta\ K_1\right). \tag{64}$$

4.2 Separation of scales: the von Neumann unitary

The commutation relations (57) show that (K_1, K_2) and (G_1, G_2) are two pairs of canonical conjugate operators. The *Stone-von Neumann uniqueness Theorem* (see [BR97] Corollary 5.2.15) assures the existence of a unitary map $\mathcal{W}$ (called *von Neumann unitary*)

$$\mathcal{W} : \mathcal{H}_{\rm phy} \longrightarrow \mathcal{H}_{\rm w} := \mathcal{H}_{\rm s} \otimes \mathcal{H}_{\rm f} = L^2(\mathbb{R}, dx_{\rm s}) \otimes L^2(\mathbb{R}, dx_{\rm f}) \tag{65}$$

such that

$$\mathcal{W} G_1 \mathcal{W}^{-1} = Q_{\rm s} = \text{multiplication by } \ x_{\rm s}, \qquad \mathcal{W} G_2 \mathcal{W}^{-1} = P_{\rm s} = -i\iota_q \delta^2 \frac{\partial}{\partial x_{\rm s}} \tag{66}$$

$$\mathcal{W} K_1 \mathcal{W}^{-1} = Q_{\rm f} = \text{multiplication by } \ x_{\rm f}, \qquad \mathcal{W} K_2 \mathcal{W}^{-1} = P_{\rm f} = -i\iota_q \ \frac{\partial}{\partial x_{\rm f}}. \tag{67}$$

The explicit costruction of the von Neumann unitary $\mathcal{W}$ is described in Appendix B.

Let $X_j := G_j + (-1)^j \delta \ K_j$ with $j = 1, 2$. From (66) and (67) it follows that

$$X_1' := \mathcal{W} X_1 \mathcal{W}^{-1} = Q_{\rm s} - \delta \ Q_{\rm f}, \qquad X_2' := \mathcal{W} X_2 \mathcal{W}^{-1} = P_{\rm s} + \delta \ P_{\rm f}. \tag{68}$$

Since X_1 and X_2 commute, one can use the spectral calculus to define any measurable function of X_1 and X_2. For any $f \in L^\infty(\mathbb{R}^2, d^2x)$ one defines $f(X_1, X_2) := \int_{\mathbb{R}^2} f(x_1, x_2) \, d\mathbb{E}^{(1)}_{x_1} \, d\mathbb{E}^{(2)}_{x_2}$ where $d\mathbb{E}^{(j)}$ is the projection-valued measure corresponding to X_j. In view of the unitarity of $\mathcal{W}$, and observing that $d\mathbb{E}'^{(j)} := \mathcal{W} d\mathbb{E}^{(j)} \mathcal{W}^{-1}$ is the projection-valued measure of X_j', one obtains that

$$\mathcal{W} f(X_1, X_2) \mathcal{W}^{-1} = \int_{\mathbb{R}^2} f(x_1, x_2) \ d{\mathbb{E}'_{x_1}}^{(1)} \ d{\mathbb{E}'_{x_2}}^{(2)} = f(X_1', X_2').$$

So the effect of the conjugation through $\mathcal{W}$ on a function f of the operators X_1 and X_2 formally amounts to replace the operators X_j with X_j' inside f.

In view of the above remark, one can easily rewrite $H^\delta_{\rm BL}$ making explicit the rôle of the fast and slow variables, obtaining

$$H^{\mathcal{W}} := \mathcal{W} H^\delta_{\rm BL} \mathcal{W}^{-1} = \mathbb{1}_{\mathcal{H}_{\rm s}} \ \otimes \ \Xi(Q_{\rm f}, P_{\rm f}) + \delta \ W(Q_{\rm f}, Q_{\rm s}, P_{\rm f}, P_{\rm s}) + \delta^2 \ V\left(P_{\rm s} + \delta \ P_{\rm f}, Q_{\rm s} - \delta \ Q_{\rm f}\right) \tag{69}$$

where, according to (61),

$$W(Q_{\rm f}, Q_{\rm s}, P_{\rm f}, P_{\rm s}) = f_1\left(P_{\rm s} + \delta \ P_{\rm f}, Q_{\rm s} - \delta \ Q_{\rm f}\right) \ Q_{\rm f} - f_2\left(P_{\rm s} + \delta \ P_{\rm f}, Q_{\rm s} - \delta \ Q_{\rm f}\right) \ P_{\rm f}. \tag{70}$$

4.3 Relevant part of the spectrum: the Landau bands

The existence of a separation between fast and slow degrees of freedom and the decomposition of the physical Hilbert space $\mathcal{H}_{\rm phy}$ into the product space $\mathcal{H}_{\rm w} = \mathcal{H}_{\rm s} \otimes \mathcal{H}_{\rm f}$ are the first two ingredients to develop the SAPT. According to the general scheme, we “replace” the canonical operators corresponding to the slow degrees of freedom with classical variables which will be re-quantized "a posteriori". Mathematically, we show that the Hamiltonian H^{W} acting in $\mathcal{H}_{\mathrm{w}}$ is the Weyl quantization of the operator-valued function (symbol) H_δ,

$$H_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) := \Xi(Q_{\mathrm{f}}, P_{\mathrm{f}}) + \delta \underbrace{W(Q_{\mathrm{f}}, x_{\mathrm{s}}, P_{\mathrm{f}}, p_{\mathrm{s}})}_{=W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})} + \delta^2 \underbrace{V\left(p_{\mathrm{s}} + \delta\, P_{\mathrm{f}}, x_{\mathrm{s}} - \delta\, Q_{\mathrm{f}}\right)}_{=V_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})}. \tag{71}$$

The quantization is defined (formally) by the rules

$$x_{\mathrm{s}} \longmapsto \mathrm{Op}_\delta(x_{\mathrm{s}}) := Q_{\mathrm{s}} \otimes \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}, \qquad p_{\mathrm{s}} \longmapsto \mathrm{Op}_\delta(p_{\mathrm{s}}) := P_{\mathrm{s}} \otimes \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}.$$

For every $(p_{\mathrm{s}}, x_{\mathrm{s}}) \in \mathbb{R}^2$, equation (71) defines an unbounded operator $H_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})$ which acts in the Hilbert space $\mathcal{H}_{\mathrm{f}}$. To make the quantization procedure rigorous, as explained in Appendix A of [PST03b], we need to consider H_δ as function from $\mathbb{R}^2$ into some Banach space which is also a domain of self-adjointness for $H_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})$. We take care of this details in the Section 4.4.

To complete the list of ingredients needed for the SAPT, we need to analize the spectrum of the principal part of the symbol (71) as $(p_{\mathrm{s}}, x_{\mathrm{s}})$ varies in $\mathbb{R}^2$. The principal part of the symbol, denoted by $H_0(p_{\mathrm{s}}, x_{\mathrm{s}})$, is given by (71) when $\delta = 0$, so it reads:

$$H_0(p_{\mathrm{s}}, x_{\mathrm{s}}) := \Xi(Q_{\mathrm{f}}, P_{\mathrm{f}}) = \frac{1}{2\Omega_\Gamma}\left[|a|^2 {P_{\mathrm{f}}}^2 + |b|^2 {Q_{\mathrm{f}}}^2 - a \cdot b\ \{Q_{\mathrm{f}}; P_{\mathrm{f}}\}\right]. \tag{72}$$

Since the principal symbol is constant on the phase space, i.e. $H_0(p_{\mathrm{s}}, x_{\mathrm{s}}) = \Xi$ for all $(p_{\mathrm{s}}, x_{\mathrm{s}}) \in \mathbb{R}^2$, we are reduced to compute the spectrum of Ξ. As well-known (see Remark 4.1 below), the spectrum of Ξ is pure point with $\sigma(\Xi) = \{\lambda_n := (n + 1/2) : n \in \mathbb{N}\}$. We refer to the eigenvalue λ_n as the n-th *Landau level.*

The spectrum of the symbol H_0 consists of a collection of constant functions $\sigma_n : \mathbb{R}^2 \to \mathbb{R}$, $n \in \mathbb{N}$, $\sigma_n(p_{\mathrm{s}}, x_{\mathrm{s}}) \equiv \lambda_n$, which we call *Landau bands.* The band σ_n is separated by the rest of the spectrum by a constant gap. In the gap condition (analogous to (31)) one can choose $C_{\mathrm{g}} = 1$. Therefore, each finite family of contiguous Landau bands defines a relevant part of the spectrun appropriate to develop the SAPT.

Remark 4.1 (The domain of self-adjointness)**.** We describe explicitly the domain of self-adjointness of $H_0(p_{\mathrm{s}}, x_{\mathrm{s}})$. Mimicking the standard theory of Landau levels, one introduces operators

$$\mathfrak{a} := \frac{i}{\sqrt{2}}\frac{\ell}{\Omega_\Gamma}\left[(a_1 + ia_2)P_{\mathrm{f}} - (b_1 + ib_2)Q_{\mathrm{f}}\right] = \frac{i}{\sqrt{2}}\left[\overline{z_a}\ P_{\mathrm{f}} - \overline{z_b}\ Q_{\mathrm{f}}\right] \tag{73}$$

$$\mathfrak{a}^\dagger := \frac{-i}{\sqrt{2}}\frac{\ell}{\Omega_\Gamma}\left[(a_1 - ia_2)P_{\mathrm{f}} - (b_1 - ib_2)Q_{\mathrm{f}}\right] = \frac{-i}{\sqrt{2}}\left[z_a\ P_{\mathrm{f}} - z_b\ Q_{\mathrm{f}}\right], \tag{74}$$

where $z_a := \frac{1}{\ell}(a_1 - ia_2)$ and $z_b := \frac{1}{\ell}(b_1 - ib_2)$. It is easy to check that

$$\mathfrak{a}\mathfrak{a}^\dagger = \Xi(Q_{\mathrm{f}}, P_{\mathrm{f}}) + \iota_q\,\frac{1}{2}\mathbb{1}_{\mathcal{H}_{\mathrm{f}}}, \qquad \mathfrak{a}^\dagger\mathfrak{a} = \Xi(Q_{\mathrm{f}}, P_{\mathrm{f}}) - \iota_q\,\frac{1}{2}\mathbb{1}_{\mathcal{H}_{\mathrm{f}}}, \qquad [\mathfrak{a}; \mathfrak{a}^\dagger] = \iota_q\ \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}. \tag{75}$$

Without loss of generality, we suppose that $\iota_q = 1$. Let ψ_0 be the *ground state* defined by $\mathfrak{a}\psi_0 = 0$. A simple computation shows that $\psi_0(x_{\mathrm{f}}) = C\mathrm{e}^{-(\beta - i\alpha)x_{\mathrm{f}}^2}$, where $C > 0$ is a



Figure 2: Structure of the spectrum of H_0. The picture shows a "relevant part of the spectrum" consisting of two Landau bands of constant energy λ_* and λ_{*+1}.

normalization constant, and $\alpha \in \mathbb{R}$, $\beta > 0$ are related to the geometry of the lattice Γ by $\alpha := {}^{a \cdot b}/_{2|a|^2}$ and $\beta := {}^{\Omega_\Gamma}/_{2|a|^2}$. Since ψ_0 is a fast decreasing smooth function, the vectors $\psi_n := (n!)^{-\frac{1}{2}}(\mathfrak{a}^\dagger)^n\psi_0$, with $n = 0, 1, \ldots$, are well defined. From the algebraic relations (75) it follows straightforwardly that: (i) $\mathfrak{a}\psi_n = \sqrt{n}\psi_{n-1}$; (ii) the family of vectors $\{\psi_n\}_{n\in\mathbb{N}}$ is an orthonormal basis for $\mathcal{H}_\mathrm{f}$ called the *generalized Hermite basis*; (iii) $\Xi\psi_n = \lambda_n\psi_n$; (iv) the spectrum of Ξ is pure point with $\sigma(\Xi) = \{\lambda_n \ : \ n \in \mathbb{N}\}$.

Let $\mathcal{L} \subset \mathcal{H}_\mathrm{f}$ be the set of the finite linear combinations of the elements of the basis $\{\psi_n\}_{n\in\mathbb{N}}$. The unbounded operators $\mathfrak{a}$, $\mathfrak{a}^\dagger$ and Ξ are well defined on $\mathcal{L}$ and on this domain $\mathfrak{a}^\dagger$ acts as the adjoint of $\mathfrak{a}$ and Ξ is symmetric. Both $\mathfrak{a}$ and $\mathfrak{a}^\dagger$ are closable and we will denote their closure by the same symbols. The operator Ξ is essentially selfadjoint on the domain $\mathcal{L}$ (the deficiency indices are both zero) and so its domain of selfadjointness $\mathcal{F} := \mathcal{D}(\Xi)$ is the closure of $\mathcal{L}$ with respect to the *graph norm* $\|\psi\|^2_\Xi := \|\psi\|^2_{\mathcal{H}_\mathrm{f}} + \|\Xi\psi\|^2_{\mathcal{H}_\mathrm{f}}$. The graph norm is equivalent to the more simple *regularized norm*

$$\|\psi\|_\mathcal{F} := \|\Xi\psi\|_{\mathcal{H}_\mathrm{f}}.$$

The domain $\mathcal{F}$ has the structure of an Hilbert space with hermitian structure provided by the *regularized scalar product* $(\psi; \varphi)_\mathcal{F} := (\Xi\psi; \Xi\varphi)_{\mathcal{H}_\mathrm{f}}$. ♦◊

4.4 Symbol class and asymptotic expansion

In this section, we firstly identify the Banach space in which the symbol H_δ defined by (71) takes values and, secondarily, we explain in which sense H_δ is a "semiclassical symbol" in a suitable Hörmander symbol class. The main results are contained in Proposition 4.2. Readers who are not interested in technical details can jump directly to the next section. For the definitions of the Hörmander classes $S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ and $S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_\mathrm{f}))$ we refer to Section 3.4.

Proposition 4.2. *Assume that* Assumption $(\mathrm{A_s})$ *holds true. Then for all* $(p_\mathrm{s},x_\mathrm{s})\in\mathbb{R}^2$ *the operator* $H_\delta(p_\mathrm{s},x_\mathrm{s})$ *is essentially self-adjoint on the dense domain* $\mathcal{L}\subset\mathcal{H}_\mathrm{f}$ *consisting of finite linear combinations of generalized Hermite functions, and its domain of self-adjointness is the domain* $\mathcal{F}$ *on which the operator* $H_0=\Xi$ *is self-adjoint. Finally,* H_δ *is in the Hörmander class* $S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_\mathrm{f}))$.

In particular, $H_\delta(p_\mathrm{s},x_\mathrm{s})$ is a bounded operator from the Hilbert space $\mathcal{F}$ to the Hilbert space $\mathcal{H}_\mathrm{f}$ for all $(p_\mathrm{s},x_\mathrm{s})\in\mathbb{R}^2$. The proof of the Proposition 4.2 follows from the *Kato-Rellich Theorem* showing that for any $(p_\mathrm{s},x_\mathrm{s})\in\mathbb{R}^2$ the operator $H_\delta(p_\mathrm{s},x_\mathrm{s})$ differs from H_0 by a relatively bounded perturbation. The latter claim will be proved in Lemmas 4.3 and 4.4 below.

In view of Assumption $(\mathrm{A_s})$, $\widetilde{\mathsf{V}}_\Gamma\in C^\infty_\mathrm{b}(\mathbb{R}^2,\mathbb{R})$ so its Fourier series

$$\widetilde{\mathsf{V}}_\Gamma(r)=\sum_{n,m\in\mathbb{Z}} w_{n,m}\ \mathrm{e}^{i2\pi n\ a^*\cdot r}\mathrm{e}^{i2\pi m\ b^*\cdot r}$$

converges uniformly and moreover $\sum_{n,m=-\infty}^{+\infty}|m|^{\alpha_1}|n|^{\alpha_2}\,|w_{n,m}|\leqslant C_\alpha$ for all $\alpha\in\mathbb{N}^2$.

Let V be the $\mathbb{Z}^2$-periodic function related to ${}^1\!/\!{\varepsilon_0}\widetilde{\mathsf{V}}_\Gamma$, as in Section 4.1. In view of (58) one has

$$\mathcal{W}\frac{\widetilde{\mathsf{V}}_\Gamma}{\mathcal{E}_0}(X_1,X_2)\mathcal{W}^{-1}=V(P_\mathrm{s}+\delta P_\mathrm{f},Q_\mathrm{s}-\delta P_\mathrm{f})=\sum_{n,m=-\infty}^{+\infty} v_{n,m}\ \mathrm{e}^{i2\pi(nP_\mathrm{s}+mQ_\mathrm{s})}\ \mathrm{e}^{i2\pi\delta(nP_\mathrm{f}-mQ_\mathrm{f})} \quad (76)$$

with $v_{n,m}:={}^1\!/\!{\varepsilon_0}w_{n,m}$ and where we used the fact that fast and slow variables commute and $[Q_\mathrm{s};P_\mathrm{s}]=\delta^2[Q_\mathrm{f};P_\mathrm{f}]$. The operator (76) can be seen as the Weyl quantization of the operator-valued symbol

$$V_\delta(p_\mathrm{s},x_\mathrm{s}):=\sum_{n,m=-\infty}^{+\infty} v_{n,m}\ \mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})}\ \mathrm{e}^{i2\pi\delta(nP_\mathrm{f}-mQ_\mathrm{f})} \quad (77)$$

with quantization rule

$$\mathrm{Op}_\delta\left(\mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})}\right)=\mathrm{e}^{i2\pi(nP_\mathrm{s}+mQ_\mathrm{s})}\otimes\mathbb{1}_{\mathcal{H}_\mathrm{f}}. \quad (78)$$

Lemma 4.3. *Let* Assumption $(\mathrm{A_s})$ *hold true. Then* $V_\delta\in S^1(\mathscr{B}(\mathcal{H}_f))\cap S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_\mathrm{f}))$. *In particular* $V_\delta(p_\mathrm{s},x_\mathrm{s})$ *is a bounded self-adjoint operator on* $\mathcal{H}_\mathrm{f}$ *for all* $(p_\mathrm{s},x_\mathrm{s})\in\mathbb{R}^2$.

Proof. It is sufficient to show that $V_\delta \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ since

$$\|\partial^\alpha V_\delta\|_{\mathscr{B}(\mathcal{F};\mathcal{H}_{\mathrm{f}})} = \|(\partial^\alpha V_\delta)\ \Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant 2\|\partial^\alpha V_\delta\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}$$

in view of $\|\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} = 2$. Let $\alpha := (\alpha_1, \alpha_2) \in \mathbb{N}^2$, then

$$\left\|\partial_{p_{\mathrm{s}}}^{\alpha_1}\partial_{x_{\mathrm{s}}}^{\alpha_2} V_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})\right\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant (2\pi)^{|\alpha|} \sum_{n,m+-\infty}^{+\infty} |n|^{\alpha_1}|m|^{\alpha_2}\,|v_{n,m}| \leqslant \frac{(2\pi)^{|\alpha|}}{\mathcal{E}_0} C_\alpha$$

for all $(p_{\mathrm{s}}, x_{\mathrm{s}}) \in \mathbb{R}^2$, as a consequence of the unitarity of $\mathrm{e}^{i2\pi\delta(nP_{\mathrm{f}}-mQ_{\mathrm{f}})}$. The self-adjointness follows by observing that $\{v_{n,m}\}$ are the Fourier coefficients of a real function. ■

Assumption $(\mathrm{A}_{\mathrm{s}})$ implies that the Γ-periodic functions $a^*\cdot\mathbf{A}_\Gamma$ and $b^*\cdot\mathbf{A}_\Gamma$ are elements of $C^\infty_{\mathrm{b}}(\mathbb{R}^2,\mathbb{R})$. By the same arguments above, one proves that the operators $f_j(P_{\mathrm{s}}+\delta P_{\mathrm{f}}, Q_{\mathrm{s}}-\delta P_{\mathrm{f}})$, $j=1,2$, appearing in (70), are the Weyl quantization of the operator-valued functions

$$f_\delta^{(j)}(p_{\mathrm{s}}, x_{\mathrm{s}}) := \sum_{n,m=-\infty}^{+\infty} f_{n,m}^{(j)}\ \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}\ \mathrm{e}^{i2\pi\delta(nP_{\mathrm{f}}-mQ_{\mathrm{f}})} \qquad j=1,2 \tag{79}$$

according to (78). The coefficients $\frac{1}{2\pi}\frac{\Phi_0}{Z\Omega_\Gamma} f_{n,m}^{(j)}$ are the Fourier coefficients of $a^*\cdot\mathbf{A}_\Gamma$ if $j=1$ and of $b^*\cdot\mathbf{A}_\Gamma$ if $j=2$. Thus, equation (70) shows that the operator $W(Q_{\mathrm{f}}, Q_{\mathrm{s}}, P_{\mathrm{f}}, P_{\mathrm{s}})$ coincides with the Weyl quantization of the operator-valued symbol

$$W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) := f_\delta^{(1)}(p_{\mathrm{s}}, x_{\mathrm{s}})\ Q_{\mathrm{f}} + f_\delta^{(2)}(p_{\mathrm{s}}, x_{\mathrm{s}})\ P_{\mathrm{f}}, \tag{80}$$

defined, initially, on the dense domain $\mathcal{L}$.

Lemma 4.4. *Let* Assumption $(\mathrm{A}_{\mathrm{s}})$ *hold true. Then* $f_\delta^{(j)} \in S^1(\mathscr{B}(\mathcal{H}_f)) \cap S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_{\mathrm{f}}))$, *for* $j=1,2$. *For all* $(p_{\mathrm{s}}, x_{\mathrm{s}}) \in \mathbb{R}^2$, *the bounded operators* $f_\delta^{(j)}(p_{\mathrm{s}}, x_{\mathrm{s}})$ *are self-adjoint while* $W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})$ *is symmetric on the dense domain* $\mathcal{L}$ *and infinitesimally bounded with respect to* Ξ. *Finally* $W_\delta \in S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_f))$.

Proof. As in the first part of the proof of Lemma 4.3, one proves that $f_\delta^{(j)} \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}})) \cap S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_{\mathrm{f}}))$ and its self-adjointness. The operator $W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})$ is a linear combination of Q_{f} and P_{f}, which are densely defined on $\mathcal{L}$, multiplied by bounded operators. Using (62) one checks by a direct computation that $W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})$ acts as a symmetric operator on $\mathcal{L}$. Since Q_{f} and P_{f} are infinitesimally bounded with respect to Ξ, then the same holds true for $W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}})$, $(p_{\mathrm{s}}, x_{\mathrm{s}}) \in \mathbb{R}^2$. The last claim follows by observing that

$$\|\partial^\alpha(f_\delta^{(j)} X_{\mathrm{f}})\|_{\mathscr{B}(\mathcal{F};\mathcal{H}_{\mathrm{f}})} = \|(\partial^\alpha f_\delta^{(j)})\ X_{\mathrm{f}}\,\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant \|X_{\mathrm{f}}\,\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}\ \|\partial^\alpha f_\delta^{(j)}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})},$$

with $j=1,2$ and $X_{\mathrm{f}} = Q_{\mathrm{f}}$ or P_{f}. Since $\|X_{\mathrm{f}}\,\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant C$ and $f_\delta^{(j)} \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$, the claim is proved. ■

Lemmas 4.3 and 4.4, together with the fact that $H_0 \equiv \Xi$ is clearly in $S^1(\mathscr{B}(\mathcal{F};\mathcal{H}_{\mathrm{f}}))$ imply the last part of Proposition 4.2.

4.5 Semiclassics: the $\mathcal{O}(\delta^4)$-approximated symbol

In this section we consider the asymptotic expansion for the symbol H_δ in the parameter δ. The Fourier expansion (77) for V_δ and the similar expression for W_δ, namely

$$W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) = \sum_{n,m=-\infty}^{+\infty} \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}\, \mathrm{e}^{i2\pi\delta(nP_{\mathrm{f}}-mQ_{\mathrm{f}})} \left[f^{(1)}_{n,m}\, Q_{\mathrm{f}} + f^{(2)}_{n,m}\, P_{\mathrm{f}}\right], \tag{81}$$

suggest a way to expand the symbol H_δ in powers of δ. By inserting the expansion $\mathrm{e}^{i2\pi\delta I_{n,m}} = \sum_{j=0}^{+\infty} \frac{(i2\pi\delta)^j}{j!} {I_{n,m}}^j$, with $I_{n,m} := nP_{\mathrm{f}} - mQ_{\mathrm{f}}$, in (77) and (81) and by exchanging the order of the series one obtains the *formal expansions*

$$V_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) \simeq \sum_{j=0}^{+\infty} \delta^j\, V_{j+2}(p_{\mathrm{s}}, x_{\mathrm{s}}) \qquad W_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) \simeq \sum_{j=0}^{+\infty} \delta^j\, W_{j+1}(p_{\mathrm{s}}, x_{\mathrm{s}}) \tag{82}$$

where

$$V_{j+2}(p_{\mathrm{s}}, x_{\mathrm{s}}) := \frac{(i2\pi)^j}{j!} \sum_{n,m=-\infty}^{+\infty} \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}\, {I_{n,m}}^j\, v_{n,m} \tag{83}$$

$$W_{j+1}(p_{\mathrm{s}}, x_{\mathrm{s}}) := \frac{(i2\pi)^j}{j!} \sum_{n,m=-\infty}^{+\infty} \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}\, {I_{n,m}}^j \left[f^{(1)}_{n,m}\, Q_{\mathrm{f}} + f^{(2)}_{n,m}\, P_{\mathrm{f}}\right]. \tag{84}$$

In view of (62), one easily shows that the operators W_j are (formally) symmetric.

The justification of the formal expansions above requires some cautions: (i) we need to specify the domains of definitions of the unbounded operators ${I_{n,m}}^j$ and consequently the domains of definitions of V_j and W_j; (ii) we need to justify the exchange of the order of the series in the equations (77) and (81).

As for (i), one notices that

$$I_{n,m} = \alpha_{n,m}\, \mathfrak{a} + \overline{\alpha_{n,m}}\, \mathfrak{a}^\dagger, \qquad \alpha_{n,m} := \frac{nz_b - mz_a}{\sqrt{2}}. \tag{85}$$

For all $(n, m) \in \mathbb{Z}^2$ the operators $I_{n,m}$ are essentially self-adjoint on the invariant dense domain $\mathcal{L}$ (their deficiency indices are both zero). The powers ${I_{n,m}}^j$ are also well defined and essentially self-adjoint on $\mathcal{L}$, as consequence of the *Nelson Theorem* (Theorem X.39 in [RS75]) since the set $\{\psi_n\}_{n\in\mathbb{N}}$ of the generalized Hermite functions is a *total set of analytic vectors* for every $I_{n,m}$ (see Example 2 Section X.6 of [RS75]). The domain of self-adjointness for ${I_{n,m}}^j$ is the closure of $\mathcal{L}$ with respect the corresponding graph norm.

The operator $V_j(p_{\mathrm{s}}, x_{\mathrm{s}})$ defined by equation (83) is an homogeneous polynomial of degree $j-2$ in $\mathfrak{a}$ and $\mathfrak{a}^\dagger$. It is symmetric (hence closable) and essentially self-adjoint on the invariant dense domain $\mathcal{L}$. Analogously, the operators

$$M^j_{n,m} := {I_{n,m}}^j \left[f^{(1)}_{n,m}\, Q_{\mathrm{f}} + f^{(2)}_{n,m}\, P_{\mathrm{f}}\right] = {I_{n,m}}^j \left[g_{n,m}\, \mathfrak{a} + \overline{g_{n,m}}\, \mathfrak{a}^\dagger\right], \qquad g_{n,m} := \frac{z_a f^{(1)}_{n,m} + z_b f^{(2)}_{n,m}}{\sqrt{2}} \tag{86}$$

which appear in the right-hand side of equation (84), are essentially self-adjoint on $\mathcal{L}$ since the set of the generalized Hermite functions provides a total set of analytic vectors. Thus we answered to point (i).

Since the generalized Hermite functions are a total set of analytic vectors for all $I_{n,m}$ then the series $\sum_{j=0}^{+\infty} \frac{(i2\pi\delta)^j}{j!} {I_{n,m}}^j \psi$ converges in norm for every $\psi \in \mathcal{L}$. From this observation, the fact that the series of coefficients $v_{n,m}$, $f^{(1)}_{n,m}$ and $f^{(2)}_{n,m}$ are absolutely convergent and that Q_f end P_f leave invariant the domain $\mathcal{L}$, one argues that for all $\psi \in \mathcal{L}$ the double series which defines $V_\delta(p_\mathrm{s}, x_\mathrm{s})\psi$ and $W_\delta(p_\mathrm{s}, x_\mathrm{s})\psi$ are absolutely convergent, hence the order of the sums can be exchanged. Thus the series appearing on the r.h.s. of (82) agrees with V_δ (respectively, with W_δ) on the dense domain $\mathcal{L}$. By a density argument, the equality in (82) holds true on the full domain of definition of V_δ (which is $\mathcal{H}_\mathrm{f}$) and W_δ respectively.

In view of the above, we write the "semiclassical expansion" of the symbol H_δ as:

$$H_\delta(p_\mathrm{s}, x_\mathrm{s}) = \Xi + \sum_{j=1}^{+\infty} \delta^j \; H_j(p_\mathrm{s}, x_\mathrm{s}), \qquad H_j(p_\mathrm{s}, x_\mathrm{s}) := W_j(p_\mathrm{s}, x_\mathrm{s}) + V_j(p_\mathrm{s}, x_\mathrm{s}) \tag{87}$$

with $V_1 = 0$.

Proposition 4.2 shows that the natural domain for the full symbol $H_\delta(p_\mathrm{s}, x_\mathrm{s})$ is the domain $\mathcal{F}$ of self-adjointness of Ξ. However, if we want to truncate the series (87) at the j-th order, we must be careful in the determination of the domain of definition of the single terms and to control the remainder. Every term in the expansion (87) is essentially self-adjoint on $\mathcal{L}$. However, the j-th order term H_j is the sum of two homogeneous polynomials in Q_f and P_f (or equivalently in $\mathfrak{a}$ and $\mathfrak{a}^\dagger$), W_j of degree j and V_j of degree $j-2$. Since $W_j = 0$ if $\mathbf{A}_\Gamma = 0$, one obtains

$$\deg\, H_j = \begin{cases} j, & \text{if } \mathbf{A}_\Gamma \neq 0 \\ j-2, & \text{if } \mathbf{A}_\Gamma = 0 \end{cases}$$

where $\deg\, H_j$ means the degree of H_j as a polynomial in Q_f and P_f. If $\deg\, H_j > 2$ then the operator H_j is not bounded by the principal symbol Ξ, and in this sense it cannot be considered as a "small perturbation" in the sense of Kato. Moreover, some other problems appear (see Remark 4.8). In order to avoid these problems, we truncate the expansion (87) up to the polynomial term of degree 2, i.e. up to order δ^2 if $\mathbf{A}_\Gamma = 0$ and up to order δ^4 if $\mathbf{A}_\Gamma \neq 0$.

Hereafter let $\natural$ be the *indicator function* of the periodic vector potential, defined as

$$\natural = \begin{cases} 0, & \text{if } A_\Gamma \neq 0 \\ 1, & \text{if } A_\Gamma = 0. \end{cases}$$

Let $\widetilde{H}^\natural_\delta(p_\mathrm{s}, x_\mathrm{s}) := \Xi + \sum_{j=1}^{2(1+\natural)} \delta^j H_j(p_\mathrm{s}, x_\mathrm{s})$, namely

$$\widetilde{H}^1_\delta(p_\mathrm{s}, x_\mathrm{s}) = \Xi + \delta^2 \sum_{n,m=-\infty}^{+\infty} v_{n,m} \; \mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})} \left(\mathbb{1}_{\mathcal{H}_\mathrm{f}} + i2\pi\delta I_{n,m} + \frac{1}{2}(i2\pi)^2\delta^2 {I_{n,m}}^2 \right) \tag{88}$$

$$\widetilde{H}^0_\delta(p_\mathrm{s}, x_\mathrm{s}) = \Xi + \delta \sum_{n,m=-\infty}^{+\infty} \mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})} \left[M^0_{n,m} + \delta(i2\pi M^1_{n,m} + v_{n,m}\mathbb{1}_{\mathcal{H}_\mathrm{f}}) \right]. \tag{89}$$

We call $\widetilde{H}^{\natural}_{\delta}$ the *approximated symbol* up to order $\delta^{2(1+\natural)}$. As a consequence of the Kato-Rellich theorem we have the following result:

Proposition 4.5. *Under* Assumption $(\mathrm{A_s})$ *there exists a constant* δ_0 *such that for every* $\delta < \delta_0$ *and for every* $(p_s, x_s) \in \mathbb{R}^2$ *the operator* $\widetilde{H}^{\natural}_{\delta}(p_s, x_s)$ *(both for* $\natural = 0$ *or 1) is self-adjoint on the domain* $\mathcal{F}$ *and bounded from below. Moreover* $\widetilde{H}^{\natural}_{\delta} \in S^1(\mathscr{B}(\mathcal{F}; \mathcal{H}_{\mathrm{f}}))$.

Proof. As proved in Lemma A.2, $\mathfrak{a}$ and $\mathfrak{a}^{\dagger}$ are infinitesimally bounded with respect to Ξ. This fact and Assumption $(\mathrm{A_s})$, which assures the fast decay of the coefficients $v_{n,m}$ and $g_{n,m}$ (see (86)), imply that the operators

$$\sum_{n,m=-\infty}^{+\infty} v_{n,m}\ \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}\left(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + i2\pi\delta I_{n,m}\right), \qquad \sum_{n,m=-\infty}^{+\infty} \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}\left[M^0_{n,m} + \delta v_{n,m}\mathbb{1}_{\mathcal{H}_{\mathrm{f}}}\right]$$

are infinitesimally bounded with respect to Ξ and are elements of $S^1(\mathscr{B}(\mathcal{F}; \mathcal{H}_{\mathrm{f}}))$.

The operators ${I_{n,m}}^2$ and $M^1_{n,m}$ are only bounded (and not infinitesimally bounded) with respect to Ξ. First of all it is easy to check that $\|{\mathfrak{a}^{\sharp}}^2\psi\|_{\mathcal{H}_{\mathrm{f}}} \leqslant 3\|\Xi\psi\|_{\mathcal{H}_{\mathrm{f}}}$ for every $\psi \in \mathcal{F}$, where $\mathfrak{a}^{\sharp}$ means $\mathfrak{a}$ or $\mathfrak{a}^{\dagger}$. Then, for every $\psi \in \mathcal{F}$,

$$\|{I_{n,m}}^2\psi\|^2_{\mathcal{H}_{\mathrm{f}}} \leqslant |\alpha_{n,m}|^4\left(\|\mathfrak{a}^2\psi\|_{\mathcal{H}_{\mathrm{f}}} + \|\{\mathfrak{a};\mathfrak{a}^{\dagger}\}^2\psi\|_{\mathcal{H}_{\mathrm{f}}} + \|{\mathfrak{a}^{\dagger}}^2\psi\|_{\mathcal{H}_{\mathrm{f}}}\right)^2 \leqslant 27\frac{d^4}{\Omega^2}(n^2+m^2)^2\ \|\Xi\psi\|^2_{\mathcal{H}_{\mathrm{f}}} \tag{90}$$

where we used the inequality $(\alpha+\beta+\gamma)^2 \leqslant 3(\alpha^2+\beta^2+\gamma^2)$, the identity $\{\mathfrak{a};\mathfrak{a}^{\dagger}\} = 2\Xi$ and the bound $|\alpha_{n,m}|^2 \leqslant {}^{d^2}\!/_{\Omega}(n^2-m^2)$ with $d^2 := \max\{|a|,|b|\}$. Assumption $(\mathrm{A_s})$ assures that the operator $\delta^4 2\pi^2\sum_{n,m=-\infty}^{+\infty} v_{n,m}\ \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})}{I_{n,m}}^2$, which appears in (88), is bounded by Ξ by a constant $\delta^4 C$, with $C \propto \sum_{n,m=-\infty}^{+\infty} v_{n,m}(n^2+m^2)$, and is in $S^1(\mathscr{B}(\mathcal{F}; \mathcal{H}_{\mathrm{f}}))$. The claim for $\widetilde{H}^1_{\delta}$ follows from the Kato-Rellich theorem fixing $\delta_0 := C^{-\frac{1}{4}}$.

The claim for $\widetilde{H}^0_{\delta}$ follows in the same way proving an inequality of the type (90) for $M^1_{n,m} = c_{n,m}\mathfrak{a}^2 + \overline{c_{n,m}}{\mathfrak{a}^{\dagger}}^2 + 2\Re(d_{n,m})\Xi + \iota_q\Im(d_{n,m})$ where $c_{n,m} := \alpha_{n,m}g_{n,m}$ and $d_{n,m} := \alpha_{n,m}\overline{g_{n,m}}$. Observe that the series of coefficients $g_{n,m}$ decays rapidly, then also the serie $c_{n,m}$ and $d_{n,m}$ have a fast decay and in particular are bounded. This implies that in the inequality of type (90) we can find a global constant which does not depend on n and m. ■

It is useful to have explicit expressions of the first terms H_j, in terms of $\mathfrak{a}$ and $\mathfrak{a}^{\dagger}$. From equations (87), (88) and (89), using the Fourier expansion of the derivatives of V and $g := {}^{1}\!/_{\sqrt{2}}\,(z_a f_1 + z_b f_2)$, it is easy to check the following:

- Case 1: $\mathbf{A}_{\Gamma} = 0$ - In this situation

$$\widetilde{H}^1_{\delta} = \Xi + \delta^2 H_2 + \delta^3 H_3 + \delta^4 H_4$$

with

$$H_2(p_{\mathrm{s}}, x_{\mathrm{s}}) = V(p_{\mathrm{s}}, x_{\mathrm{s}})\ \mathbb{1}_{\mathcal{H}_{\mathrm{f}}} \tag{91}$$

$$H_3(p_{\mathrm{s}}, x_{\mathrm{s}}) = -\frac{1}{\sqrt{2}}\left[D_z(V)\ \mathfrak{a} + D_{\overline{z}}(V)\ \mathfrak{a}^{\dagger}\right] \tag{92}$$

$$H_4(p_{\mathrm{s}}, x_{\mathrm{s}}) = \frac{1}{4}\left[|D_z|^2(V)\ 2\Xi + D^2_z(V)\ \mathfrak{a}^2 + D^2_{\overline{z}}(V)\ {\mathfrak{a}^{\dagger}}^2\right] \tag{93}$$

where D_z is the differential operator defined by $D_z := \left(z_a \frac{\partial}{\partial x_{\mathrm{s}}} - z_b \frac{\partial}{\partial p_{\mathrm{s}}}\right)$ and $D_{\overline{z}}$ is obtained by replacing z_a and z_b with $\overline{z}_a$ and $\overline{z}_b$. Since V is real, $D_{\overline{z}}(V) = \overline{D_z(V)}$, which shows that H_3 is symmetric. The explicit expression of the second order differential operator $|D_z|^2 := D_z \circ D_{\overline{z}}$ is

$$|D_z|^2 = \frac{1}{\Omega_\Gamma}\left(|a|^2 \, \frac{\partial^2}{\partial x_{\mathrm{s}}^2} - 2a \cdot b \, \frac{\partial^2}{\partial x_{\mathrm{s}} \partial p_{\mathrm{s}}} + |b|^2 \, \frac{\partial^2}{\partial p_{\mathrm{s}}^2}\right). \tag{94}$$

- Case 2: $\mathsf{A}_\Gamma \neq 0$ - In this situation

$$\widetilde{H}^0_\delta = \Xi + \delta H_1 + \delta^2 H_2$$

with

$$H_1(p_{\mathrm{s}}, x_{\mathrm{s}}) = g(p_{\mathrm{s}}, x_{\mathrm{s}}) \ \mathfrak{a} + \overline{g}(p_{\mathrm{s}}, x_{\mathrm{s}}) \ \mathfrak{a}^\dagger \tag{95}$$

$$H_2(p_{\mathrm{s}}, x_{\mathrm{s}}) = V \ \mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \sqrt{2} D_z(\overline{g}) \ \Xi - \frac{1}{\sqrt{2}}\left[D_z(g) \ \mathfrak{a}^2 + D_{\overline{z}}(\overline{g}) \ {\mathfrak{a}^\dagger}^2\right]. \tag{96}$$

In the computation of (96) we used the first of the gauge conditions (52) which assures that

$$D_z(\overline{g}) = \frac{1}{\sqrt{2}\Omega_\Gamma}\left[|a|^2 \, \frac{\partial f_1}{\partial x_{\mathrm{s}}} + a \cdot b \left(\frac{\partial f_2}{\partial x_{\mathrm{s}}} - \frac{\partial f_1}{\partial p_{\mathrm{s}}}\right) - |b|^2 \, \frac{\partial f_2}{\partial p_{\mathrm{s}}}\right]$$

is a real function. From the definition of g, f_1 and f_2 it follows that

$$g(a^* \cdot r, b^* \cdot r) = \pi \sqrt{2} \frac{Z \ell}{\Phi_0} \left[(\mathsf{A}_\Gamma)_1 - i(\mathsf{A}_\Gamma)_2\right](r), \tag{97}$$

namely g is the dimensionless $\mathbb{Z}^2$-periodic function related to the Γ-periodic function $(\mathsf{A}_\Gamma)_1 - i(\mathsf{A}_\Gamma)_2$, up to a multiplicative constant.

4.6 Main result: effective dynamics for strong magnetic fields

Preliminary estimates on the remainder

The difference $\mathfrak{R}^\natural_\delta := H_\delta - \widetilde{H}^\natural_\delta$ is a self-adjoint element of $S^1(\mathscr{B}(\mathcal{F}; \mathcal{H}_{\mathrm{f}}))$, which we call the *remainder symbol.* To develop the SAPT for the Harper regime we need to estimate the order of the remainder symbol. The next result shows essentially that

$$H_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) = \widetilde{H}^\natural_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) + \mathcal{O}(\delta^{2(\natural+1)}), \qquad \pi_{\mathrm{r}} \ H_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) \ \pi_{\mathrm{r}} = \pi_{\mathrm{r}} \ \widetilde{H}^\natural_\delta(p_{\mathrm{s}}, x_{\mathrm{s}}) \ \pi_{\mathrm{r}} + \mathcal{O}(\delta^{2(\natural+1)+1}), \tag{98}$$

where

$$\pi_{\mathrm{r}} := \sum_{i=1}^{m} |\psi_{k_i}\rangle\langle\psi_{k_i}| \tag{99}$$

is the projection on the subspace spanned by the finite family of generalized Hermite functions $\{\psi_{k_i}\}_{i=1}^m$. In other words, the error done by replacing the true symbol H_δ with the approximated symbol $\widetilde{H}^{\natural}_{\delta}$ (which has order $2(\natural+1)$ in δ) is of the same order of the approximated symbol, so in this sense $\widetilde{H}^{\natural}_{\delta}$ is not a good approximation for H_{δ}. On the other side, what we need to develop the SAPT is to control the operator $\pi_{\mathrm{r}}\ H_{\delta}\ \pi_{\mathrm{r}}$, which is well approximated by $\pi_{\mathrm{r}}\ \widetilde{H}^{\natural}_{\delta}\ \pi_{\mathrm{r}}$ up to an error of order $2(\natural+1)+1$ in δ.

Proposition 4.6. *Let* Assumption ($\mathrm{A_s}$) *hold true. Then* $\mathfrak{R}^{\natural}_{\delta}$ *has order* $\mathcal{O}(\delta^{2(\natural+1)})$*, i. e. there exist a constant* C *such that* $\|\mathfrak{R}^{\natural}_{\delta}(p_s,x_s)\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} \leqslant C\delta^{2(\natural+1)}$ *for all* $(p_s,x_s)\in\mathbb{R}^2$. *Moreover* $\|\mathfrak{R}^{\natural}_{\delta}\ \pi_r\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} = \|\pi_r\ \mathfrak{R}^{\natural}_{\delta}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant C\delta^{2(\natural+1)+1}$*, for all* $(p_s,x_s)\in\mathbb{R}^2$*, i.e.* $\mathfrak{R}^{\natural}_{\delta}\ \pi_r$*,* $\pi_r\ \mathfrak{R}^{\natural}_{\delta}$ *and* $[\mathfrak{R}^{\natural}_{\delta};\pi_r]$ *are* $\mathscr{B}(\mathcal{H}_{\mathrm{f}})$*-valued symbols of order* $\mathcal{O}(\delta^{2(\natural+1)+1})$.

Proof. **(Case $\natural=1$)** The explicit expression of the remainder symbol is

$$\mathfrak{R}^{1}_{\delta}(p_{\mathrm{s}},x_{\mathrm{s}}) = \delta^2 \sum_{n,m=-\infty}^{+\infty} v_{n,m}\ \mathrm{e}^{i2\pi(np_{\mathrm{s}}+mx_{\mathrm{s}})} \left[\mathrm{e}^{i2\pi\delta I_{n,m}} - \left(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + i2\pi\delta I_{n,m} + \frac{1}{2}(i2\pi)^2\delta^2 {I_{n,m}}^2\right)\right]. \tag{100}$$

and from (100) it follows that $\|\mathfrak{R}^{1}_{\delta}(p_{\mathrm{s}},x_{\mathrm{s}})\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} \leqslant \delta^2\sum_{n,m=-\infty}^{+\infty}|v_{n,m}|\Lambda_{n,m}$ with

$$\Lambda_{n,m} := \sup_{\psi\in\mathcal{H}_{\mathrm{f}}}\setminus\{0\}\frac{\left\|\left[\mathrm{e}^{i2\pi\delta I_{n,m}} - \left(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + i2\pi\delta I_{n,m} + \frac{1}{2}(i2\pi)^2\delta^2 {I_{n,m}}^2\right)\right]\Xi^{-1}\psi\right\|_{\mathcal{H}_{\mathrm{f}}}}{\|\psi\|_{\mathcal{H}_{\mathrm{f}}}} \tag{101}$$

since $\|\psi\|_{\mathcal{F}} := \|\Xi\psi\|_{\mathcal{H}_{\mathrm{f}}}$ and $\mathcal{F}=\Xi^{-1}\mathcal{H}_{\mathrm{f}}$. The operators $I_{n,m}$ are essentially self-adjoint on $\mathcal{L}$ and we denote their closure with the same symbol. Since the operators ${I_{n,m}}^2$ are positive, we can consider the resolvent operators $R_{n,m} := ({I_{n,m}}^2+\mathbb{1}_{\mathcal{H}_{\mathrm{f}}})^{-1}$. Let suppose that

$$\zeta_{n,m}(\delta) := \left\|\left[\mathrm{e}^{i2\pi\delta I_{n,m}} - \left(\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + i2\pi\delta I_{n,m} + \frac{1}{2}(i2\pi)^2\delta^2 {I_{n,m}}^2\right)\right] R_{n,m}\right\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant \zeta(\delta), \tag{102}$$

for all $n,m\in\mathbb{Z}$, with $\sup_{\delta}\zeta(\delta)<+\infty$. Then equation (101) would imply

$$\Lambda_{n,m} \leqslant \zeta(\delta)\left\|({I_{n,m}}^2+\mathbb{1}_{\mathcal{H}_{\mathrm{f}}})\,\Xi^{-1}\right\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}.$$

Noticing that ${I_{n,m}}^2 = {\alpha_{n,m}}^2\ \mathfrak{a}^2 + \overline{\alpha_{n,m}}^2\ {\mathfrak{a}^{\dagger}}^2 + 2|\alpha_{n,m}|^2\ \Xi$ and observing that $\|\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}=2$, $\|\mathfrak{a}^2\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}=1$ and $\|{\mathfrak{a}^{\dagger}}^2\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})}=2\sqrt{2}$, one deduces from inequality (102) that

$$\|\mathfrak{R}^{1}_{\delta}(p_{\mathrm{s}},x_{\mathrm{s}})\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_{\mathrm{f}})} \leqslant C_1\left(\sum_{n,m=-\infty}^{+\infty}|v_{n,m}|(|n|+|m|)^2\right)\delta^2\zeta(\delta) \leqslant C_2\delta^2\zeta(\delta)$$

for suitable positive constants C_1 and C_2.

It remains to prove the inequality (102) and the estimate on $\zeta(\delta)$. By spectral calculus one has that $\zeta_{n,m}(\delta)=\sup_{t\in\sigma(I_{n,m})}|Z_{\delta}(t)| \leqslant \sup_{t\in\mathbb{R}}|Z_{\delta}(t)| =: \zeta(\delta)$ where

$$Z_{\delta}(t) := 4\pi^2\delta^2\ \frac{\mathrm{e}^{i2\pi\delta t} - \left(1 + i2\pi\delta t - \frac{1}{2}(2\pi\delta t)^2\right)}{(2\pi\delta t)^2 + 4\pi^2\delta^2}.$$

After some manipulations and the change of variable $\tau := 2\pi\delta t$ one has that

$$G_\delta(\tau) := \frac{1}{4\pi^4\delta^4}\left|Z_\delta\left(\frac{\tau}{2\pi\delta}\right)\right|^2 \leqslant \frac{\tau^4 + 4\tau^2\cos(\tau) - 8\tau\sin(\tau) - 8\cos(\tau) + 8}{\tau^4} < C_3.$$

Thus $\zeta(\delta)^2 = 4\pi^4\delta^4\ \sup_{\tau\in\mathbb{R}} G_\delta(\tau) \leqslant 4\pi^4 C_3\delta^4$, hence $\|\mathfrak{R}^1_\delta(p_\mathrm{s}, x_\mathrm{s})\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_\mathrm{f})} \leqslant C\delta^4$. This concludes the first part of the proof.

Since $\|\mathfrak{R}^1_\delta\pi_\mathrm{r}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant \sum_{i=1}^m \|\mathfrak{R}^1_\delta|\psi_{k_i}\rangle\langle\psi_{k_i}|\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$, then it is enough to show that for any Hermite vector ψ_k the inequality $\|\mathfrak{R}^1_\delta|\psi_k\rangle\langle\psi_k|\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C_k\delta^5$ holds true. Observing that $\|\mathfrak{R}^1_\delta|\psi_k\rangle\langle\psi_k|\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = \|\mathfrak{R}^1_\delta\psi_k\|_{\mathcal{H}_\mathrm{f}}$, one deduces

$$\lim_{\delta\to 0}\delta^{-5}\|\mathfrak{R}^1_\delta|\psi_k\rangle\langle\psi_k|\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = \lim_{\delta\to 0}\left\|\sum_{n,m=-\infty}^{+\infty} v_{n,m}\mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})}\left(\sum_{j=3}^{+\infty}\frac{(i2\pi)^j\delta^{j-3}}{j!}{I_{n,m}}^j\right)\psi_k\right\|_{\mathcal{H}_\mathrm{f}}$$

$$\leqslant \frac{4}{3}\pi^3\sum_{n,m=-\infty}^{+\infty}|v_{n,m}|\ \|{I_{n,m}}^3\psi_k\|_{\mathcal{H}_\mathrm{f}} \leqslant \frac{32}{3}\pi^3 C'\|{A^\dagger}^3\psi_k\|_{\mathcal{H}_\mathrm{f}} = \frac{32}{3}\sqrt{(k+3)!}\ \pi^3 C' =: C_k$$

where $C' := \sum_{n,m=-\infty}^{+\infty}|v_{n,m}|\ |\alpha_{n,m}|^3$ is finite in view of Assumption ($\mathrm{A_s}$).

This shows that for all $\delta \in [0,\delta_0)$ (for a suitable $\delta_0 > 0$) the norm $\|\mathfrak{R}^1_\delta|\psi_k\rangle\langle\psi_k|\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$ is bounded by $C_k\delta^5$ and so it follows that $\|\mathfrak{R}^1_\delta\pi_\mathrm{r}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant mC\delta^5$ with $C := \max_{1,\ldots,m}\{C_{k_i}\}$. Finally $\|\pi_\mathrm{r}\mathfrak{R}^1_\delta\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = \|(\mathfrak{R}^1_\delta\pi_\mathrm{r})^\dagger\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} = \|\mathfrak{R}^1_\delta\pi_\mathrm{r}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$.

(Case $\natural = 0$) The proof proceeds as in the previous case. Divide the remainder symbol in two terms $\mathfrak{R}^0_\delta = \mathfrak{R}^0_0 + \mathfrak{R}^0_1$ where:

$$\mathfrak{R}^0_0(p_\mathrm{s}, x_\mathrm{s}) := \delta\sum_{n,m=-\infty}^{+\infty}\mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})}\left(\mathrm{e}^{i2\pi\delta I_{n,m}} - \mathbb{1}_{\mathcal{H}_\mathrm{f}} - i2\pi\delta I_{n,m}\right)M^0_{n,m}$$

$$\mathfrak{R}^0_1(p_\mathrm{s}, x_\mathrm{s}) := \delta^2\sum_{n,m=-\infty}^{+\infty} v_{n,m}\ \mathrm{e}^{i2\pi(np_\mathrm{s}+mx_\mathrm{s})}\left(\mathrm{e}^{i2\pi\delta I_{n,m}} - \mathbb{1}_{\mathcal{H}_\mathrm{f}}\right).$$

The control of $\mathfrak{R}^0_1$ is easy, indeed $\|\mathfrak{R}^0_1\|_{\mathscr{B}(\mathcal{D},\mathcal{H}_\mathrm{f})} \leqslant 2\|\mathfrak{R}^0_1\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant 4C\delta^2$ where $C := \sum_{n,m=-\infty}^{+\infty}|v_{n,m}|$. Moreover (with the same technique used for the case $\natural = 1$), one can check that for any Hermite vector ψ_k the function $t_1(\delta) := \frac{1}{\delta^3}\|\mathfrak{R}^0_1(p_\mathrm{s}, x_\mathrm{s})\psi_k\|_{\mathcal{H}_\mathrm{f}}$ is bounded by a constant $C_k > 0$ in a suitable interval $[0,\delta_0)$. This assures that $\|\mathfrak{R}^0_1\ \pi_\mathrm{r}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$ is of order $\mathcal{O}(\delta^3)$.

To control $\mathfrak{R}^0_0$ we need to estimate $\Sigma_{n,m} := \|\left(\mathrm{e}^{i2\pi\delta I_{n,m}} - \mathbb{1}_{\mathcal{H}_\mathrm{f}} - i2\pi\delta I_{n,m}\right)M^0_{n,m}\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$. Let $R'_{n,m}$ be the resolvent $(I_{n,m}+i\mathbb{1}_{\mathcal{H}_\mathrm{f}})^{-1}$. It is easy to check that $\|(I_{n,m}+i\mathbb{1}_{\mathcal{H}_\mathrm{f}})M^0_{n,m}\Xi^{-1}\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$ is bounded by a linear expression in $|n|$ and $|m|$. Indeed, as proved in Proposition 4.5, both $M^0_{n,m}$ and $M^1_{n,m}$ are bounded by Ξ. By spectral calculus $\|\left(\mathrm{e}^{i2\pi\delta I_{n,m}} - \mathbb{1}_{\mathcal{H}_\mathrm{f}} - i2\pi\delta I_{n,m}\right)R'_{n,m}\|^2_{\mathscr{B}(\mathcal{H}_\mathrm{f})}$ is bounded by the maximum in τ of the function $F_\delta(\tau) := 4\pi^2\delta^2\ \frac{\tau^2-2\tau\sin\tau-2\cos\tau+2}{\tau^2}$.

The last part follows observing that $M^0_{n,m}$ is a linear combinations of $\mathfrak{a}$ and $\mathfrak{a}^\dagger$ and so they act splitting a Hermite vector ψ_k as $c^k_{n,m}\psi_{k-1} + d^k_{n,m}\psi_{k+1}$ where for a fixed k the coefficients depend on $f^{(j)}_{n,m}$. To conclude the proof it is sufficient to notice that $t_0(\delta) := \frac{1}{\delta^2}\|\left(\mathrm{e}^{i2\pi\delta I_{n,m}} - \mathbb{1}_{\mathcal{H}_\mathrm{f}} - i2\pi\delta I_{n,m}\right)\psi_k\|_{\mathcal{H}_\mathrm{f}}$ is bounded by a constant $C_k > 0$ in a suitable interval $[0,\delta_0)$. ■

Derivation of the adiabatically decoupled effective dynamics

We recall that the Weyl quantization of the symbol H_δ is the Hamiltonian (69), namely $\mathrm{Op}_\delta(H_\delta) = H^{\mathcal{W}}$. As for the approximated symbol $\widetilde{H}^{\natural}_{\delta}$, we pose $\widetilde{H}^{\natural} := \mathrm{Op}_\delta(\widetilde{H}^{\natural}_{\delta})$. Both $H^{\mathcal{W}}$ and $\widetilde{H}^{\natural}$ are bounded operators from $L^2(\mathbb{R}, dx_{\mathrm{s}}) \otimes \mathcal{F}$ to $\mathcal{H}_{\mathrm{w}} := L^2(\mathbb{R}, dx_{\mathrm{s}}) \otimes \mathcal{H}_{\mathrm{f}}$.

Theorem 4.7. *Let* Assumption ($\mathrm{A_s}$) *be satisfied. Let* $\{\sigma_n(\cdot)\}_{n\in\mathcal{I}}$, *with* $\mathcal{I} = \{n, \ldots, n+m-1\}$, *be a family of Landau bands for* Ξ *and let* $\pi_{\mathrm{r}} := \sum_{n\in\mathcal{I}} |\psi_n\rangle\langle\psi_n|$ *be the spectral projector of* $H_0 = \Xi$ *corresponding to the set* $\{\sigma_n(p_{\mathrm{s}}, x_{\mathrm{s}})\}_{n\in\mathcal{I}}$. *Then:*

1. Almost-invariant subspace: *there exists an orthogonal projection* $\Pi^{\natural}_{\delta} \in \mathscr{B}(\mathcal{H}_{\mathrm{w}})$, *with* $\Pi^{\natural}_{\delta} = \mathrm{Op}_\delta(\pi) + \mathcal{O}_0(\delta^\infty)$, $\pi(p_{\mathrm{s}}, x_{\mathrm{s}}) \asymp \sum_{j=0}^{\infty} \delta^j\ \pi_j(p_{\mathrm{s}}, x_{\mathrm{s}})$, *and* $\pi_0(p_{\mathrm{s}}, x_{\mathrm{s}}) \equiv \pi_{\mathrm{r}}$, *such that*

$$[\widetilde{H}^{\natural};\Pi^{\natural}_{\delta}] = \mathcal{O}_0(\delta^\infty), \qquad [H^{\mathcal{W}};\Pi^{\natural}_{\delta}] = \mathcal{O}_0(\delta^{2(\natural+1)+1}). \tag{103}$$

2. Effective dynamics: *let* $\Pi_{\mathrm{r}} := \mathbb{1}_{\mathcal{H}_{\mathrm{s}}} \otimes \pi_{\mathrm{r}} \in \mathscr{B}(\mathcal{H}_{\mathrm{w}})$ *and* $\mathcal{K} := \mathrm{Ran}\ \Pi_{\mathrm{r}} \simeq L^2(\mathbb{R}, dx_s) \otimes \mathbb{C}^m$. *Then there exists a unitary operator* $U^{\natural}_{\delta} \in \mathscr{B}(\mathcal{H}_{\mathrm{w}})$ *such that*

- (i) $U^{\natural}_{\delta} = \mathrm{Op}_\delta(u) + \mathcal{O}_0(\delta^\infty)$, *where the symbol* $u \asymp \sum_{j=0}^{\infty} \delta^j u_j$ *has principal part* $u_0 \equiv \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}$;
- (ii) $\Pi_{\mathrm{r}} = U^{\natural}_{\delta}\ \Pi^{\natural}_{\delta}\ {U^{\natural}_{\delta}}^{-1}$;
- (iii) *Let* $h^{\natural}$ *in* $S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ *be a resummation of the formal symbol* $u \,\sharp\, \pi \,\sharp\, \widetilde{H}^{\natural}_{\delta} \,\sharp\, \pi \,\sharp\, u^\dagger$ *and define the* effective Hamiltonian *by* $H^{\delta}_{\mathrm{eff}} := \mathrm{Op}_\delta(h^{\natural})$. *Since* $[H^{\delta}_{\mathrm{eff}}; \Pi_{\mathrm{r}}] = 0$, $H^{\delta}_{\mathrm{eff}}$ *is a bounded operator on* $\mathcal{K}$. *Then*

$$U^{\natural}_{\delta}\, \Pi^{\natural}_{\delta}\, H^{\mathcal{W}}\, \Pi^{\natural}_{\delta}\, {U^{\natural}_{\delta}}^{-1} = H^{\delta}_{\mathrm{eff}} + \mathcal{O}_0(\delta^{2(\natural+1)+1}) \in \mathscr{B}(\mathcal{K}). \tag{104}$$

2'. Effective dynamics for a single Landau band when $\mathbf{A}_\Gamma = 0$***:*** *Consider a single Landau band* $\sigma_*(\cdot) = \lambda_*$, *so that* $\pi_{\mathrm{r}} = |\psi_*\rangle\langle\psi_*|$. *Then, up to the order* δ^4, *one has that*

$$H^{\delta}_{\mathrm{eff}} = \lambda_* \mathbb{1}_{\mathcal{H}_{\mathrm{s}}} + \delta^2\ V(P_{\mathrm{s}}, Q_{\mathrm{s}}) + \delta^4\ \frac{\lambda_*}{2}\ Y(P_{\mathrm{s}}, Q_{\mathrm{s}}) + \mathcal{O}_0\left(\delta^5\right) \tag{105}$$

where $V(P_{\mathrm{s}}, Q_{\mathrm{s}}) := \mathrm{Op}_\delta(V)$ *is the Weyl quantization of the* $\mathbb{Z}^2$*-periodic function* $V(p_{\mathrm{s}}, x_{\mathrm{s}})$ *related to the* Γ*-periodic potential* V_Γ, *while* $Y(P_{\mathrm{s}}, Q_{\mathrm{s}}) := \mathrm{Op}_\delta(|D_z|^2(V))$ *is the Weyl quantization of the function* $|D_z|^2(V)(p_{\mathrm{s}}, x_{\mathrm{s}})$ *defined through the differential operator* (94).

The derivation of the effective dynamics when $\mathbf{A}_\Gamma \neq 0$ will be considered in Section 4.8.

Proof of Theorem 4.7

Step 1. Almost-invariant subspace

As explained in the first part of proof of the Theorem 3.12 one constructs a formal symbol π (the *Moyal projection*) such that: (i) $\pi \,\sharp\, \pi \asymp \pi$; (ii) $\pi^\dagger = \pi$; (iii) $\widetilde{H}^{\natural}_{\delta} \,\sharp\, \pi \asymp \pi \,\sharp\, \widetilde{H}^{\natural}_{\delta}$. Such a symbol $\pi \asymp \sum_{j=0}^{\infty} \delta^j \pi_j$ is constructed recursively order by order starting from $\pi_0 = \pi_{\mathrm{r}}$ and $\widetilde{H}^{\natural}_{\delta}$ and it is unique (see Lemma 2.3. in [PST03b]). The recursive relations are

$$\pi_n := \pi_n^{\mathrm{D}} + \pi_n^{\mathrm{OD}} \tag{106}$$

where the *diagonal part* is $\pi_n^{\mathrm{D}} := \pi_{\mathrm{r}} G_n \pi_{\mathrm{r}} + (\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \pi_{\mathrm{r}}) G_n (\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \pi_{\mathrm{r}})$ with

$$G_n := \left[\left(\sum_{j=0}^{n-1} \delta^j \pi_j \right) \sharp \left(\sum_{j=0}^{n-1} \delta^j \pi_j \right) - \left(\sum_{j=0}^{n-1} \delta^j \pi_j \right) \right]_n . \tag{107}$$

The *off-diagonal part* is defined by the implicit relation $[H_0; \pi_n^{\mathrm{OD}}] = -F_n$ where

$$F_n := \left[\widetilde{H}^{\natural}_{\delta} \sharp \left(\sum_{j=0}^{n-1} \delta^j \pi_j + \delta^n \pi_n^{\mathrm{D}} \right) - \left(\sum_{j=0}^{n-1} \delta^j \pi_j + \delta^n \pi_n^{\mathrm{D}} \right) \sharp \widetilde{H}^{\natural}_{\delta} \right]_n . \tag{108}$$

The uniqueness allows us to construct π only locally and this local construction is explained in the second part of Lemma 2.3 in [PST03b]. In our case we can choose a $(p_{\mathrm{s}}, x_{\mathrm{s}})$-independent positively oriented complex circle $\Lambda \subset \mathbb{C}$, symmetric with respect to the real axis, which encloses the family of (constant) spectral bands $\{\sigma_n(\cdot) = \lambda_n\}_{n \in \mathcal{I}}$ and such that $\mathrm{dist}(\Lambda, \sigma(H_0)) \geqslant \frac{1}{2}$ (see Figure 2). For all $\lambda \in \Lambda$ we construct recursively the *Moyal resolvent* (or *parametrix*) $R^{\natural}(\lambda; \cdot) := \sum_{j=0}^{\infty} \delta^j R^{\natural}_j(\lambda; \cdot)$ of $\widetilde{H}^{\natural}_{\delta}$, following the same technique explained during the proof of Theorem 3.12. The approximants of the symbol π are related to the approximants of the Moyal resolvent by the usual Riesz formula $\pi_j(z) := \dfrac{i}{2\pi} \oint_{\Lambda} d\lambda \; R^{\natural}_j(\lambda; z)$ where $z := (p_{\mathrm{s}}, x_{\mathrm{s}}) \in \mathbb{R}^2$. Some care is required to show (iii) since, by construction, $\widetilde{H}^{\natural}_{\delta} \sharp \pi$ takes values in $\mathscr{B}(\mathcal{H}_{\mathrm{f}})$ while $\pi \sharp \widetilde{H}^{\natural}_{\delta}$ takes values in $\mathscr{B}(\mathcal{F})$. To solve this problem one can use the same argument proposed in Lemma 7 of [PST03a].

The technical and new part of the proof consist in showing that $\pi \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}})) \cap S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}, \mathcal{F}))$. The Riesz formula implies $\|(\partial^{\alpha}_{z} \pi_j)(z)\|_{\flat} \leqslant 2\pi \; \sup_{\lambda \in \Lambda} \|\partial^{\alpha}_{z} R^{\natural}_j(\lambda; z)\|_{\flat}$ for all $\alpha \in \mathbb{N}^2$ ($\flat$ means either $\mathscr{B}(\mathcal{H}_{\mathrm{f}})$ or $\mathscr{B}(\mathcal{H}_{\mathrm{f}}; \mathcal{F})$ and $\partial^{\alpha}_{z} := \partial^{\alpha_1}_{p_{\mathrm{s}}} \partial^{\alpha_2}_{x_{\mathrm{s}}}$) since Λ does not depend on z. Then we need only to show that $R^{\natural}_j(\lambda; \cdot) \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}})) \cap S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}, \mathcal{F}))$. The choice of Λ assures $\|R^{\natural}_0(\lambda; z)\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} = \|(\Xi - \lambda \mathbb{1}_{\mathcal{H}_{\mathrm{f}}})^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant 2$. Moreover $\partial^{\alpha}_{z} R^{\natural}_0(\lambda; z) = 0$ for all $\alpha \neq 0$ and this implies that $R^{\natural}_0 \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ uniformly in λ. Since $\|R^{\natural}_0(\lambda; z)\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}}, \mathcal{F})} = \|\Xi(\Xi - \lambda \mathbb{1}_{\mathcal{H}_{\mathrm{f}}})^{-1}\|_{\mathscr{B}(\mathcal{H}_{\mathrm{f}})} \leqslant \infty$ one concludes that $R^{\natural}_0 \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}, \mathcal{F}))$ uniformly in λ.

By means of equation (43), one has $R^{\natural}_j = -R^{\natural}_0 L^{\natural}_j$ where $L^{\natural}_j$ is the *j-th order obstruction* for $R^{\natural}_0$ to be the Moyal resolvent. In view of this recursive relation, the proof of $R^{\natural}_j \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ for all $j \in \mathbb{N}$ is reduced to show that $L^{\natural}_j \in S^1(\mathscr{B}(\mathcal{H}_{\mathrm{f}}))$ for all $j \in \mathbb{N}$.

The first order obstruction, computed by means of (44), is

$$L^{\natural}_1(\lambda; z) = \delta^{-1}[(\widetilde{H}^{\natural}_{\delta}(z) - \lambda \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}) \sharp R^{\natural}_0(\lambda; z) - \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}]_1 = H_1(z) \; R^{\natural}_0(\lambda; z) - \frac{i}{2} \{\Xi; R^{\natural}_0(\lambda; z)\}_{p_{\mathrm{s}}, x_{\mathrm{s}}}.$$

Since Ξ and $R^{\natural}_0$ do not depend on $z \in \mathbb{R}^2$ it follows that $L^{\natural}_1 = H_1 R^{\natural}_0$. The operator H_1 is linear in $\mathfrak{a}$ and $\mathfrak{a}^{\dagger}$ (with all its derivative) if $\natural = 0$ or $H_1 = 0$ if $\natural = 1$. In both cases H_1 (with its derivatives) is infinitesimally bounded with respect to Ξ (see Lemma A.2). This shows that $L_1^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ (but not in $S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}, \mathcal{F}))$ if $\natural = 0$).

We proceed by induction assuming that $L_j^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ for all $j \leqslant m \in \mathbb{N}$. The $(m+1)$-th order obstruction $L_{m+1}^\natural$ can be computed by means of equation (44) and the Moyal formula for the expansion of $\sharp$ (see equation (A.9) in [Teu03]). After some manipulations, one gets

$$L_{m+1}^\natural(\lambda; z) = \frac{1}{(2i)^{m+1}} \sum_{\substack{\alpha_1+\alpha_2+r+l=m+1 \\ 0\leqslant l \leqslant m,\ 0\leqslant r\leqslant 2(\natural+1)}} \frac{(-1)^{|\alpha|+1}}{\alpha_1!\alpha_2!} \left(\partial_{x_\mathrm{s}}^{\alpha_1} \partial_{p_\mathrm{s}}^{\alpha_2} H_r R_0^\natural\right)(\lambda; z) \ \left(\partial_{p_\mathrm{s}}^{\alpha_1} \partial_{x_\mathrm{s}}^{\alpha_2} L_l^\natural\right)(\lambda; z).$$

Since $H_r R_0^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ uniformly in λ (see Proposition 4.6) then $L_{m+1}^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$, and this concludes the inductive argument.

Finally to prove $R_j^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}, \mathcal{F}))$, observe that $\|\partial_z^\alpha R_j^\natural\|_{\mathscr{B}(\mathcal{H}_\mathrm{f},\mathcal{F})} = \|\Xi\, R_0^\natural\, (\partial_z^\alpha L_j^\natural)\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant C_\alpha \|\Xi\, R_0^\natural\|_{\mathscr{B}(\mathcal{H}_\mathrm{f})} \leqslant +\infty$ for all $j, \alpha \in \mathbb{N}$.

Remark 4.8. It clearly emerges from the proof that the order $\delta^{2(\natural+1)}$ is the best approximation which can be obtained with this technique. The obstruction is the condition $H_r R_0^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$, which can be satisfied by the resolvent $R_0^\natural := (\Xi - \zeta \mathbb{1}_\mathrm{f})^{-1}$ only for $0 \leqslant r \leqslant 2(\natural + 1)$. ♦◊

Proposition A.9 of [Teu03] assures that $\widetilde{H}_\delta^\natural \,\sharp\, \pi \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$ and, by adjointness, also $\pi \,\sharp\, \widetilde{H}_\delta^\natural \in S^1(\mathscr{B}(\mathcal{H}_\mathrm{f}))$. By construction $[\widetilde{H}^\natural; \mathrm{Op}_\delta(\pi)] = \mathrm{Op}_\delta([\widetilde{H}_\delta^\natural; \pi]_\sharp) = \mathcal{O}_0(\delta^\infty)$ where $[\widetilde{H}_\delta^\natural; \pi]_\sharp := \widetilde{H}_\delta^\natural \,\sharp\, \pi - \pi \,\sharp\, \widetilde{H}_\delta^\natural = \mathcal{O}(\delta^\infty)$ denotes the *Moyal commutator*. Observing that $[H_\delta; \pi]_\sharp = [\widetilde{H}_\delta^\natural + \mathfrak{R}_\delta^\natural; \pi]_\sharp = [\mathfrak{R}_\delta^\natural; \pi]_\sharp + \mathcal{O}(\delta^\infty)$ and since Proposition 4.6 implies $[\mathfrak{R}_\delta^\natural; \pi]_\sharp = [\mathfrak{R}_\delta^\natural; \pi_\mathrm{r}] + \mathcal{O}(\delta^{2(\natural+1)+1}) = \mathcal{O}(\delta^{2(\natural+1)+1})$, it follows $[H_\delta; \pi]_\sharp = \mathcal{O}(\delta^{2(\natural+1)+1})$ which implies after the quantization $[H^\mathcal{W}; \mathrm{Op}_\delta(\pi)] = \mathcal{O}_0(\delta^{2(\natural+1)+1})$.

The last step is to obtain the true projection $\Pi_\delta^\natural$ (the *super adiabatic projection*) from $\mathrm{Op}_\delta(\pi)$ by means of the formula (46). Since $\Pi_\delta^\natural - \mathrm{Op}_\delta(\pi) = \mathcal{O}_0(\delta^\infty)$, one recovers the estimates (103).

Step 2. Construction of the intertwining unitary

The construction of the intertwining unitary follows as in the proof of Theorem 3.1 of [PST03b]. Firstly one constructs a formal symbol $u \asymp \sum_{j=0}^\infty \delta^j u_j$ such that: (i) $u^\dagger \,\sharp\, u = u \,\sharp\, u^\dagger = \mathbb{1}_{\mathcal{H}_\mathrm{f}}$; (ii) $u \,\sharp\, \pi \,\sharp\, u^\dagger = \pi_\mathrm{r}$.

The existence of such a symbol follows from a recursive procedure starting from u_0 (which can be fixed to be $\mathbb{1}_{\mathcal{H}_\mathrm{f}}$ in our specific case) and using the expansion of $\pi \asymp \sum_{j=0}^\infty \delta^j \pi_j$ obtained above. However, the symbol u which comes out of this procedure is not unique. The recursive relations are

$$u_n := a_n + b_n \qquad \text{with} \qquad a_n := -\frac{1}{2} A_n, \qquad b_n := [\pi_\mathrm{r}; B_n] \tag{109}$$

where

$$A_n := \left[\left(\sum_{j=0}^{n-1} \delta^j u_j \right) \sharp \left(\sum_{j=0}^{n-1} \delta^j u_j \right)^\dagger - \mathbb{1}_{\mathcal{H}_{\rm f}} \right]_n \tag{110}$$

and

$$B_n := \left[\left(\sum_{j=0}^{n-1} \delta^j u_j + \delta^n a_n \right) \sharp\, \pi \,\sharp \left(\sum_{j=0}^{n-1} \delta^j u_j + \delta^n a_n \right)^\dagger - \pi_{\rm r} \right]_n \tag{111}$$

Since $u_0 = \mathbb{1}_{\mathcal{H}_{\rm f}} \in S^1(\mathscr{B}(\mathcal{H}_{\rm f}))$, then it follows by induction that $u_j \in S^1(\mathscr{B}(\mathcal{H}_{\rm f}))$ for all $j \in \mathbb{N}$.

The quantization of u is an element of $\mathscr{B}(\mathcal{H}_{\rm w})$ but it is not a true unitary. Nevertheless $\mathrm{Op}_\delta(u)$ can be modified by an $\mathcal{O}_0(\delta^\infty)$ term using the same technique of Lemma 3.3 (Step II) in [PST03b] to obtain the true unitary $U^\natural_\delta$.

Step 3. Effective dynamics

By construction $[H^\delta_{\rm eff}; \Pi_{\rm r}] = \mathrm{Op}_\delta([h^\natural; \pi_{\rm r}]_\sharp) = [U^\natural_\delta\, \Pi^\natural_\delta\, \widetilde{H}^\natural_\delta\, \Pi^\natural_\delta\, {U^\natural_\delta}^{-1}; \Pi_{\rm r}] = 0$ since $\Pi_{\rm r} = U^\natural_\delta\, \Pi^\natural_\delta\, {U^\natural_\delta}^{-1}$. Moreover equation (104) follows observing that $U^\natural_\delta\, \Pi^\natural_\delta\, H^{\mathcal{W}}\, \Pi^\natural_\delta\, {U^\natural_\delta}^{-1} - H^\delta_{\rm eff}$ coincides with the quantization of $u \,\sharp\, \pi \,\sharp\, \mathfrak{R}^\natural_\delta \,\sharp\, \pi \,\sharp\, u^\dagger$ which is a symbol of order $\mathcal{O}(\delta^{2(\natural+1)+1})$.

Step 4. The case of a single Landau band when $\mathbf{A}_\Gamma = 0$

We need to expand the Moyal product $h^{\natural=1} = u \,\sharp\, \pi \,\sharp\, \widetilde{H}^1_\delta \,\sharp\, \pi \,\sharp\, u^\dagger = \pi_{\rm r} \,\sharp\, u \,\sharp\, \widetilde{H}^1_\delta \,\sharp\, u^\dagger \,\sharp\, \pi_{\rm r} + \mathcal{O}(\delta^\infty)$ up to the order δ^4. To compute the various terms of the expansion $h^{\natural=1} \asymp \sum_{j=0}^\infty \delta^j h_j$ it is useful to define $\chi_j := [u \,\sharp\, \widetilde{H}^1_\delta \,\sharp\, u^\dagger]_j$, so that $h_j = \pi_{\rm r} \chi_j \pi_{\rm r}$. Observing that

$$u \,\sharp\, \widetilde{H}^1_\delta - \left(\sum_{j=0}^{m-1} \delta^j\ \chi_j \right) \sharp\, u = \left(u \,\sharp\, \widetilde{H}^1_\delta \,\sharp\, u^\dagger - \sum_{j=0}^{m-1} \delta^j \chi_j \right) \sharp\, u + \mathcal{O}(\delta^\infty) = \delta^m \chi_m + \mathcal{O}(\delta^{m+1})$$

one obtains the useful formula

$$\chi_m = \left[u \,\sharp\, \widetilde{H}^1_\delta - \left(\sum_{j=0}^{m-1} \delta^j\ \chi_j \right) \sharp\, u \right]_m . \tag{112}$$

At the zeroth order one finds $h_0 = \pi_0\, u_0\, H_0\, u_0^\dagger\, \pi_0 = \pi_{\rm r}\, \Xi\, \pi_{\rm r} = \lambda_*\, \pi_{\rm r}$ since $u_0 = \mathbb{1}_{\mathcal{H}_{\rm f}}$ and $\pi_0 = \pi_{\rm r}$. Its quantization is the operator $\mathrm{Op}_\delta(h_0) = \lambda_* \mathbb{1}_{\mathcal{H}_{\rm s}}$ acting on $\mathcal{K} = L^2(\mathbb{R}, dx_{\rm s})$.

As for the first order ($m = 1$), $\chi_1 = u_1 H_0 + u_0 H_1 - \chi_0 u_1 + [u_0 \,\sharp\, H_0]_1 - [\chi_0 \,\sharp\, u_0]_1 = [u_1; \Xi]$ since $\chi_0 = u_0 H_0 u_0^{-1} = \Xi$ and $H_1 = 0$. Then $h_1 = \pi_{\rm r}[u_1; \Xi]\pi_{\rm r} = \lambda_*(\pi_{\rm r} u_1 \pi_{\rm r} - \pi_{\rm r} u_1 \pi_{\rm r}) = 0$, hence $\mathrm{Op}_\delta(h_1) = 0$.

At the second order ($m = 2$), one obtains after some manipulations $\chi_2 = H_2 + u_2\, \Xi - \Xi\, u_2 - \chi_1\, u_1$ which implies $h_2 = \pi_{\rm r} H_2 \pi_{\rm r} - \pi_{\rm r} \chi_1 u_1 \pi_{\rm r}$. We need to compute u_1. Using equations (109), (110) and (111) one obtains that $-2a_1 := \left[u_0 \,\sharp\, u_0{}^\dagger - \mathbb{1}_{\mathcal{H}_{\rm f}}\right]_1 = 0$ and $b_1 := [\pi_{\rm r}; B_1]$ with $B_1 = [u_0 \,\sharp\, \pi \,\sharp\, u_0{}^* - \pi_{\rm r}]_1 = \pi_1$ since $a_1 = 0$. To compute π_1 we use equations (106), (107) and

(108). Since $G_1 = [\pi_{\mathrm{r}} \sharp \pi_{\mathrm{r}} - \pi_{\mathrm{r}}]_1 = 0$ it follows that $\pi_1^{\mathrm{D}} = 0$. In the case of a single energy band in the relevant part of the spectrum, the implicit relation which defines π_n^{OD} can be solved, obtaining the useful equation

$$\pi_n^{\mathrm{OD}} = \pi_{\mathrm{r}} F_n (\Xi - \lambda_* \mathbb{1}_{\mathcal{H}_{\mathrm{f}}})^{-1} (\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \pi_{\mathrm{r}}) - (\mathbb{1}_{\mathcal{H}_{\mathrm{f}}} - \pi_{\mathrm{r}}) (\Xi - \lambda_* \mathbb{1}_{\mathcal{H}_{\mathrm{f}}})^{-1} F_n \pi_{\mathrm{r}}. \tag{113}$$

Since $F_1 = [\widetilde{H}^1_\delta \sharp \pi_{\mathrm{r}} - \pi_{\mathrm{r}} \sharp \widetilde{H}^1_\delta]_1 = H_1 \pi_{\mathrm{r}} - \pi_{\mathrm{r}} H_1 = 0$, being $H_1 = 0$, it follows $B_1 = \pi_1 = \pi_1^{\mathrm{OD}} = 0$ and consecutively $u_1 = b_1 = 0$. Then $h_2 = \pi_{\mathrm{r}} \, H_2 \, \pi_{\mathrm{r}} = V \, \pi_{\mathrm{r}}$, according to (91), and its quantization defines on $\mathcal{K}$ the operator $\mathrm{Op}_\delta(h_2) = V(P_{\mathrm{s}}, X_{\mathrm{s}})$.

Considering (112) at the third order (m=3) and using $u_1 = 0$, one obtains after some computations $\chi_3 = H_3 + u_3 \, \Xi - \Xi \, u_3 - \chi_1 \, u_2$ which implies $h_3 = \pi_{\mathrm{r}} \, H_3 \, \pi_{\mathrm{r}} - \pi_{\mathrm{r}} \, \chi_1 \, u_2 \, \pi_{\mathrm{r}}$. Thus we need to compute u_2. Since $u_1 = 0$, it follows $-2a_2 = [u_0 \sharp u_0{}^\dagger - \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}]_2 = 0$, $B_2 = [u_0 \sharp \pi \sharp u_0{}^\dagger - \pi_{\mathrm{r}}]_2 = \pi_2$ and $b_2 = [\pi_{\mathrm{r}}; \pi_2]$. Since $\pi_1 = 0$, one has that $G_2 = [\pi_{\mathrm{r}} \sharp \pi_{\mathrm{r}} - \pi_{\mathrm{r}}]_2 = 0$ which implies $\pi_2^{\mathrm{D}} = 0$. To compute π_2^{OD} we need $F_2 = [\widetilde{H}^1_\delta \sharp \pi_{\mathrm{r}} - \pi_{\mathrm{r}} \sharp \widetilde{H}^1_\delta]_2 = [H_2; \pi_{\mathrm{r}}] = [\mathbb{1}_{\mathcal{H}_{\mathrm{f}}}; \pi_{\mathrm{r}}] = 0$, where $H_2 = V \, \mathbb{1}_{\mathcal{H}_{\mathrm{f}}}$ has been used. Then $B_2 = \pi_2 = \pi_2^{\mathrm{OD}} = 0$ and consequently $u_2 = b_2 = 0$. Therefore $h_3 = \pi_{\mathrm{r}} H_3 \pi_{\mathrm{r}}$, and equation (92) implies that $\pi_{\mathrm{r}} H_3 \pi_{\mathrm{r}} = 0$ in view of $\pi_{\mathrm{r}} \mathfrak{a} \pi_{\mathrm{r}} = \langle \psi_* | \mathfrak{a} | \psi_* \rangle \pi_{\mathrm{r}} = 0$ and similarly for $\mathfrak{a}^\dagger$. Then $\mathrm{Op}_\delta(h_3) = 0$.

To compute the fourth order, we do not need to compute u_3 and π_3. Indeed, by computing (112) at the fourth order (m=4) one finds $\chi_4 = H_4 + u_4 \, \Xi - \Xi \, u_4 + u_3 \, H_1 - \chi_3 \, u_1 = H_4 + u_4 \, \Xi - \Xi \, u_4$ since $H_1 = u_1 = u_2 = 0$. Then $h_4 = \pi_{\mathrm{r}} H_4 \pi_{\mathrm{r}} = \frac{\lambda_*}{2} |D_z(V)|^2 \, \pi_{\mathrm{r}}$, according to equation (93), and its quantization yields $\mathrm{Op}_\delta(h_4) = \frac{\lambda_*}{2} Y(P_{\mathrm{s}}, Q_{\mathrm{s}})$.

4.7 Harper-like Hamiltonians

The first term in (105) is a multiple of the identity, and therefore does not contribute to the dynamics as far as the expectation values of the observables are concerned. The leading term, providing a non-trivial contribution to the dynamics at the original microscopic time scale $s \propto \delta^2 \tau$, is the bounded operator $V(P_{\mathrm{s}}, Q_{\mathrm{s}})$ acting on the reference Hilbert space $L^2(\mathbb{R}, dx_{\mathrm{s}})$. This operator is the Weyl quantization of the $\mathbb{Z}^2$-periodic smooth function V defined on the classical phase space $\mathbb{R}^2$. Hereafter we write $x_{\mathrm{s}} \equiv x$ to simplify the notation.

The quantization procedure can be reformulated by introducing the unitary operators $\mathcal{U}_\infty := \mathrm{e}^{-i2\pi Q_{\mathrm{s}}}$ and $\mathcal{V}_\infty := \mathrm{e}^{-i2\pi P_{\mathrm{s}}}$ (*Harper unitaries*), acting on $L^2(\mathbb{R}, dx)$ as (recall $\delta^2 = {}^{h_B}\!/\!{}_{2\pi}$)

$$(\mathcal{U}_\infty \psi)(x) = \mathrm{e}^{-i2\pi x} \psi(x), \qquad\qquad (\mathcal{V}_\infty \psi)(x) = \psi(x - \iota_q h_B). \tag{114}$$

For any $\mathbb{Z}^2$-periodic function $F(p, x) = \sum_{n,m=-\infty}^{+\infty} f_{n,m} \mathrm{e}^{-i2\pi(np+mx)}$ whose Fourier series is uniformly convergent, the h_B-Weyl quantization of F is given by the formula

$$\widehat{F}(\mathcal{U}_\infty, \mathcal{V}_\infty) = \sum_{n,m=-\infty}^{+\infty} f_{n,m} \mathrm{e}^{-i\pi nm(\iota_q h_B)} \, \mathcal{V}_\infty^n \mathcal{U}_\infty^m. \tag{115}$$

where the fundamental commutation relation $\mathcal{U}_\infty \mathcal{V}_\infty = \mathrm{e}^{-i2\pi(\iota_q h_B)} \, \mathcal{V}_\infty \mathcal{U}_\infty$ has been used. Formula (115) defines a *Harper-like Hamiltonian* with deformation parameter h_B. Indeed, the special case $H_{\mathrm{Har}} = \mathcal{U}_\infty + \mathcal{U}_\infty^{-1} + \mathcal{V}_\infty + \mathcal{V}_\infty^{-1}$ is the celebrated Harper Hamiltonian, namely the operator acting on $L^2(\mathbb{R}, dx)$ as

$$(H_{\mathrm{Har}}\psi)(x) = \psi(x + h_B) + \psi(x - h_B) + 2\cos(2\pi x)\psi(x). \tag{116}$$

In analogy with Section 3, we summarize the discussion in the following conclusion

Conclusion 4.9. *Under the assumptions of Theorem 4.7, for every* $\mathsf{V}_\Gamma \in C_{\mathrm{b}}^\infty(\mathbb{R}^2, \mathbb{R})$*, in the Harper regime* ($h_B \to 0$)*, the dynamics generated by the Hamiltonian* $\mathbf{H}_{\mathrm{BL}}$ (1) *restricted to the spectral subspace corresponding to a Landau level* λ_* *is approximated up to an error of order* h_B *(and up to a unitary transform and an energy rescaling) by the dynamics generated on the reference Hilbert space* $L^2(\mathbb{R}, dx)$ *by a Harper-like Hamiltonian, i.e. by a power series in the Harper unitaries* $\mathcal{U}_\infty$ *and* $\mathcal{V}_\infty$*, defined by* (114).

4.8 Coupling of Landau bands in a periodic magnetic potential

According to Theorem 4.7, the first non-trivial term which describes the effective dynamics in the almost invariant subspace related to a single Landau band λ_* is of order $\delta^2 \propto h_B$. An important ingredient in the proof is that $\mathbf{A}_\Gamma = 0$ implies $H_1 = 0$. Moreover, the second non-trivial correction appears at order $\delta^4 \propto {h_B}^2$ although $H_3 \neq 0$. Indeed, the correction at order δ^3 vanishes since H_3, defined by (92), is linear in $\mathfrak{a}$ and $\mathfrak{a}^\dagger$, hence $\langle\psi_*|H_3|\psi_*\rangle = 0$. This observation suggests that for a family of Landau bands which contains two contiguous bands $\{\lambda_*, \lambda_{*+1}\}$ one has, in general, a second non-trivial correction of order $\delta^3 \propto {h_B}^{\frac{3}{2}}$ for the effective dynamics. Indeed, in this case one has $\pi_{\mathrm{r}} H_3 \pi_{\mathrm{r}} \neq 0$ since $\langle\psi_*|H_3|\psi_{*+1}\rangle$ is generally non zero. Nevertheless, also in this case, the first non-trivial correction is of order δ^2.

Is there any mechanism to produce a non-trivial correction in the effective dynamics with leading order $\delta \propto {h_B}^{\frac{1}{2}}$? An affirmative answer requires $H_1 \neq 0$, and the latter condition is satisfied if we include in the Hamiltonian $\mathbf{H}_{\mathrm{BL}}$ the effect of a Γ-periodic vector potential $\mathbf{A}_\Gamma$ (i.e. $\natural = 0$). Since in this situation H_1 is linear in $\mathfrak{a}$ and $\mathfrak{a}^\dagger$, to obtain a non-trivial effect we need to consider a spectral subspace which contains at least two contiguous Landau bands.

Our goal is to derive the (non-trivial) leading order for the effective Hamiltonian in this framework. According to the notation of Theorem 4.7, we need to expand the Moyal product $h^{\natural=0} = u \sharp \pi \sharp \widetilde{H}_\delta^0 \sharp \pi \sharp u^\dagger = \pi_{\mathrm{r}} \sharp u \sharp \widetilde{H}_\delta^0 \sharp u^\dagger \sharp \pi_{\mathrm{r}} + \mathcal{O}(\delta^\infty)$ up to the first order δ. The symbols $\pi = \pi_{\mathrm{r}} + \mathcal{O}(\delta)$ and $u = \mathbb{1}_{\mathcal{H}_{\mathrm{f}}} + \mathcal{O}(\delta)$ are derived as in the general construction showed in the proof of Theorem 4.7. Now $\mathcal{K} := \mathrm{Ran}\ \Pi_{\mathrm{r}} \simeq L^2(\mathbb{R}, dx_{\mathrm{s}}) \otimes \mathbb{C}^2$.

Expanding at zero order one finds $h_0 = \pi_0\, u_0\, H_0 u_0^{-1}\, \pi_0 = \pi_{\mathrm{r}}\, \Xi\, \pi_{\mathrm{r}} = \pi_{\mathrm{r}}\, \Xi = \Xi\, \pi_{\mathrm{r}}$ and its quantization is the operator on $\mathcal{K}$ defined by

$$\mathrm{Op}_\delta(h_0) = \begin{pmatrix} \left(n_* + \frac{3}{2}\right) \mathbb{1}_{\mathcal{H}_{\mathrm{s}}} & 0 \\ 0 & \left(n_* + \frac{1}{2}\right) \mathbb{1}_{\mathcal{H}_{\mathrm{s}}} \end{pmatrix} = (n_* + 1)\mathbb{1}_{\mathcal{K}} + \begin{pmatrix} \frac{1}{2}\mathbb{1}_{\mathcal{H}_{\mathrm{s}}} & 0 \\ 0 & -\frac{1}{2}\mathbb{1}_{\mathcal{H}_{\mathrm{s}}} \end{pmatrix} \tag{117}$$

As for the next order, from equation (112) it follows $\chi_1 = H_1 + u_1 H_0 - H_0 u_1$ (we use $\chi_0 = H_0$) which implies $h_1 = \pi_{\mathrm{r}} \chi_1 \pi_{\mathrm{r}} = \pi_{\mathrm{r}}\, H_1\, \pi_{\mathrm{r}} + \pi_{\mathrm{r}}\, [u_1; \Xi]\, \pi_{\mathrm{r}}$. To conclude the computation we need u_1 and π_1. Using the recursive formulas (106), (107), (108), (109), (110) and (111), one obtains $-2a_1 := \left[u_0 \sharp u_0{}^\dagger - \mathbb{1}_{\mathcal{H}_\mathrm{f}}\right]_1 = 0$, $b_1 := [\pi_\mathrm{r}; B_1]$ and $B_1 = [u_0 \sharp \pi \sharp u_0{}^* - \pi_\mathrm{r}]_1 = \pi_1$ since $a_1 = 0$. Observing that $G_1 = [\pi_\mathrm{r} \sharp \pi_\mathrm{r} - \pi_\mathrm{r}]_1 = 0$, it follows that $\pi_1^\mathrm{D} = 0$ and so $u_1 = [\pi_\mathrm{r}; \pi_1] = [\pi_\mathrm{r}; \pi_1^\mathrm{OD}]$ which implies $\pi_\mathrm{r} u_1 \pi_\mathrm{r} = 0$. Finally $\pi_\mathrm{r}[u_1; \Xi]\pi_\mathrm{r} = \pi_\mathrm{r}[u_1; \pi_\mathrm{r}\Xi\pi_\mathrm{r}]\pi_\mathrm{r} = 0$ and so $h_1 = \pi_\mathrm{r} H_1 \pi_\mathrm{r}$. According to (95) the quantization of h_1 is an operator which acts on $\mathcal{K}$ as

$$\mathrm{Op}_\delta(h_1) = \sqrt{n_* + 1} \begin{pmatrix} 0 & \mathcal{G}(P_\mathrm{s}, Q_\mathrm{s}) \\ & \\ \mathcal{G}(P_\mathrm{s}, Q_\mathrm{s})^\dagger & 0 \end{pmatrix} \tag{118}$$

where the operator $\mathcal{G}(P_\mathrm{s}, Q_\mathrm{s})$ is defined on $L^2(\mathbb{R}, dx_\mathrm{s})$ as the Weyl quantization of the $\mathbb{Z}^2$-periodic function g defined by equation (97). Summarizing, we obtained the following result:

Theorem 4.10 (Effective Hamiltonian with a periodic magnetic potential)**.** *Under the assumptions of Theorem 4.7, in the case* $\mathbf{A}_\Gamma \neq 0$ *the dynamics in the spectral subspace related to a family of two contiguous Landau bands* $\{\sigma_{*+j}(\cdot) = \lambda_{*+j} \mid j = 0, 1\}$ *is approximated by the effective Hamiltonian* $H^\delta_\mathrm{eff} := Op_\delta(h^{(\natural=0)})$ *on the reference space* $\mathcal{K} = L^2(\mathbb{R}, dx_s) \otimes \mathbb{C}^2$ *which is given, up to errors of order* δ^2, *by*

$$H^\delta_\mathrm{eff} = (n_* + 1)\mathbb{1}_\mathcal{K} + \sqrt{n_* + 1} \begin{pmatrix} \frac{1}{2\sqrt{n_*+1}}\mathbb{1}_{\mathcal{H}_s} & \delta\ \mathcal{G}(P_s, Q_s) \\ & \\ \delta\ \mathcal{G}(P_s, Q_s)^\dagger & -\frac{1}{2\sqrt{n_*+1}}\mathbb{1}_{\mathcal{H}_s} \end{pmatrix} + \mathcal{O}_0\left(\delta^2\right), \tag{119}$$

according to the notation introduced in (117) *and* (118)*.*

Equation (97) shows that $g(p_\mathrm{s}, x_\mathrm{s}) = g_1(p_\mathrm{s}, x_\mathrm{s}) - i g_2(p_\mathrm{s}, x_\mathrm{s})$ where the function g_1 and g_2 are related to the component $(\mathsf{A}_\Gamma)_1$ and $(\mathsf{A}_\Gamma)_2$ of the Γ-periodic vector potential by the relation $g_j(a^* \cdot r, b^* \cdot r) = \pi\sqrt{2}\frac{Z\ell}{\Phi_0}(\mathsf{A}_\Gamma)_j(r)$, $j = 1, 2$. Let $\mathcal{G}_j(P_\mathrm{s}, Q_\mathrm{s})$ be the Weyl quantization of g_j. By introducing the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \qquad \sigma_\perp = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \tag{120}$$

one can rewrite the effective Hamiltonian (119) in the form

$$H^\delta_\mathrm{eff} = \left((n_* + 1)\mathbb{1}_{\mathbb{C}^2} + \frac{1}{2}\sigma_\perp\right) \otimes \mathbb{1}_{\mathcal{H}_\mathrm{s}} + \delta\sqrt{n_* + 1} \ \sum_{j=1}^{2} \sigma_j \otimes \mathcal{G}_j(P_\mathrm{s}, Q_\mathrm{s}) + \mathcal{O}_0\left(\delta^2\right). \tag{121}$$

Clearly, the operator $\mathcal{G}_j(P_\mathrm{s}, Q_\mathrm{s})$ are Harper-like Hamiltonians and can be represented as a power series of the Harper unitaries $\mathcal{U}_\infty$ and $\mathcal{V}_\infty$ of type (115). In this case the coefficients in the expansion are (up to a multiplicative constant) the Fourier coefficients of the components $(\mathsf{A}_\Gamma)_j$ of the Γ-periodic vector potential.

The determination of the spectrum of H^δ_eff can be reduced to the (generally simpler) problem of the computation of the spectrum of $\mathcal{G}\mathcal{G}^\dagger$.

Proposition 4.11. *Let* $H^{\delta=1}_\mathrm{eff}$ *be the first order approximation of the effective Hamiltonian* (119) *(or* (121)*). Then*

$$\sigma(H^{\delta=1}_\mathrm{eff}) = (n_* + 1) + S_+ \cup S_-, \qquad S_\pm := \{\pm\sqrt{1/4 + \delta^2(n_* + 1)\ \lambda}\ :\ \lambda \in \overline{\sigma}(\mathcal{G}\mathcal{G}^\dagger)\}$$

where $\overline{\sigma}(\mathcal{G}\mathcal{G}^\dagger) = \sigma(\mathcal{G}\mathcal{G}^\dagger) \cup \{0\}$ *if* $\{0\} \in \sigma(\mathcal{G}^\dagger\mathcal{G}) \setminus \sigma(\mathcal{G}\mathcal{G}^\dagger)$ *and* $\overline{\sigma}(\mathcal{G}\mathcal{G}^\dagger) = \sigma(\mathcal{G}\mathcal{G}^\dagger)$ *otherwise.*

Proof. We give only a sketch of the proof. The term $(n_* + 1)\mathbb{1}_{\mathcal{K}}$ shifts the spectrum of a constant value $(n_* + 1)$, then we can consider only the spectrum of $\mathcal{B} := H^{\delta=1}_{\text{eff}} - (n_* + 1)\mathbb{1}_{\mathcal{K}}$. A simple computation shows that

$$\mathcal{B}^2 = \begin{pmatrix} \frac{1}{2}\mathbb{1}_{\mathcal{H}_\text{s}} & \delta\sqrt{n_* + 1}\ \mathcal{G} \\ \delta\sqrt{n_* + 1}\ \mathcal{G}^\dagger & -\frac{1}{2}\mathbb{1}_{\mathcal{H}_\text{s}} \end{pmatrix}^2 = \frac{1}{4}\mathbb{1}_{\mathcal{K}} + \delta^2(n_* + 1)\begin{pmatrix} \mathcal{G}\mathcal{G}^\dagger & 0 \\ 0 & \mathcal{G}^\dagger\mathcal{G} \end{pmatrix}$$

which implies that $\sigma(\mathcal{B}^2) = \{1/4 + \delta^2(n_* + 1)\lambda \ : \ \lambda \in \sigma(\mathcal{G}\mathcal{G}^\dagger) \cup \sigma(\mathcal{G}^\dagger\mathcal{G})\}$. The operators $\mathcal{G}\mathcal{G}^\dagger$, $\mathcal{G}^\dagger\mathcal{G}$ and $\mathcal{B}$ are bounded and self-adjoint. To show that $\sigma(\mathcal{G}\mathcal{G}^\dagger) \setminus \{0\} = \sigma(\mathcal{G}^\dagger\mathcal{G}) \setminus \{0\}$, let $\lambda \in \sigma(\mathcal{G}\mathcal{G}^\dagger)$ with $\lambda \neq 0$ and $\{\psi_n\}_{n\in\mathbb{N}} \subset \mathcal{H}_\text{s} \setminus \text{Ker}(\mathcal{G}^\dagger)$ be a sequence of non zero vectors such that $\|(\mathcal{G}\mathcal{G}^\dagger - \lambda)\psi_n\|_{\mathcal{H}_\text{s}} \to 0$ (*Weyl's criterion*), then $\|(\mathcal{G}^\dagger\mathcal{G} - \lambda)\mathcal{G}^\dagger\psi_n\|_{\mathcal{H}_\text{s}} \leqslant \|\mathcal{G}^\dagger\|_{\mathscr{B}(\mathcal{H}_\text{s})}\|(\mathcal{G}\mathcal{G}^\dagger - \lambda)\psi_n\|_{\mathcal{H}_\text{s}} \to 0$. This implies that $\sigma(\mathcal{G}\mathcal{G}^\dagger) \cup \sigma(\mathcal{G}^\dagger\mathcal{G}) = \overline{\sigma}(\mathcal{G}\mathcal{G}^\dagger)$. Now let $\varepsilon_\pm(\lambda) := \pm\sqrt{1/4 + \delta^2(n_* + 1)\ \lambda}$ with $\lambda \in \overline{\sigma}(\mathcal{G}\mathcal{G}^\dagger)$ and $\{\psi_n\}_{n\in\mathbb{N}}$ a sequence of generalized eigenvectors for $\mathcal{G}\mathcal{G}^\dagger$ relative to λ. Then $\Psi^{(\pm)}_n := ((1/2 + \varepsilon_\pm)\psi_n, \delta\sqrt{n_* + 1}\mathcal{G}^\dagger\psi_n) \in \mathcal{H}_\text{s} \otimes \mathbb{C}^2$ is a sequence of generalized eigenvectors for $\mathcal{B}$ relative to $\varepsilon_\pm$. ■

A Some technical results

Since we include in our analysis a periodic vector potential A_Γ (as a new ingredient with respect to the standard literature) we include a short discussion of the self-adjointness and the spectral properties of the operators H_{BL} and H_{per}.

A.1 Self-adjointness of H_{BL} and H_{per}

The *second Sobolev space* $\mathcal{H}^2(\mathbb{R}^2)$ is defined to be the set of all $\psi \in L^2(\mathbb{R}^2)$ such that $\partial^{n_1}_{x_1}\partial^{n_2}_{x_2}\psi \in L^2(\mathbb{R}^2)$ in the sense of distributions for all $n := (n_1, n_2) \in \mathbb{N}^2$ with $|n| := n_1 + n_2 \leqslant 2$. One can proves that $\mathcal{H}^2(\mathbb{R}^2)$ is the closure of $C^\infty_\text{c}(\mathbb{R}^2, \mathbb{C})$ with respect to the *Sobolev norm* $\|\cdot\|_{\mathcal{H}^2} := \|(\mathbb{1} - \Delta_x)\cdot\|_{L^2}$ and has a Hilbert space structure. Similarly the *second magnetic-Sobolev space* $\mathcal{H}^2_\text{M}(\mathbb{R}^2)$ is defined to be the set of all $\psi \in L^2(\mathbb{R}^2)$ such that $D^{n_1}_1 D^{n_2}_2\psi \in L^2(\mathbb{R}^2)$ in the sense of distributions for all $n \in \mathbb{N}^2$ with $|\alpha| \leqslant 2$, where $D_1 := (\partial_{x_1} + \frac{i}{2}x_2)$ and $D_2 := (\partial_{x_2} - \frac{i}{2}x_1)$. One can prove that $\mathcal{H}^2_\text{M}(\mathbb{R}^2)$ is the closure of $C^\infty_\text{c}(\mathbb{R}^2, \mathbb{C})$ with respect to the *magnetic-Sobolev norm* $\|\cdot\|_{\mathcal{H}^2_\text{M}} := \|(\mathbb{1} - \Delta_M)\cdot\|_{L^2}$, where $\Delta_M := {D_1}^2 + {D_2}^2$ is the *magnetic-Laplacian.* Moreover, $\mathcal{H}^2_\text{M}(\mathbb{R}^2)$ has a natural Hilbert space structure. For further details see Section IX.6 and IX.7 of [RS75] and Chapter 7 of [LL01].

Proof of Proposition 2.1.
We prove the claim for the dimensionless operators, namely we fix all the physical constants equal to 1 in (9) and (15).

- *Step 1.* First of all we prove that H_{per} is essentially self-adjoint on $C^\infty_\text{c}(\mathbb{R}^2, \mathbb{C})$ and self-adjoint on $\mathcal{H}^2(\mathbb{R}^2)$. Notice that

$$H_{\text{per}} = \frac{1}{2}\left[-i\nabla_x - A_\Gamma(x)\right]^2 + V_\Gamma(x) = -\frac{1}{2}\Delta_x + T_1 + \frac{1}{2}T_2 \tag{122}$$

with $T_1 := iA_\Gamma \cdot \nabla_x$ and $T_2 := i(\nabla_x \cdot A_\Gamma) + |A_\Gamma|^2 + 2V_\Gamma$. The free Hamiltonian $-{}^1\!/\!{}_2\Delta_x$ is a self-adjoint operator with domain $\mathcal{H}^2(\mathbb{R}^2)$, essentially self-adjoint on $C^\infty_{\mathrm{c}}(\mathbb{R}^2, \mathbb{C})$ and from Assumption ($\mathrm{A_w}$) it follows that T_2 is an infinitesimally bounded with respect to $-{}^1\!/\!{}_2\Delta_x$ (notice that $T_2 - 2V_\Gamma$ is bounded). The symmetric operator T_1 is unbounded with domain $\mathcal{D}(T_1) \supset \mathcal{H}^2(\mathbb{R}^2)$. Let $\psi \in \mathcal{H}^2(\mathbb{R}^2)$, then

$$\left\|(A_\Gamma)_j \partial_{x_j} \psi\right\|^2_{L^2} \leqslant \|(A_\Gamma)_j\|^2_\infty \int_{\mathbb{R}^2} \lambda_j^2 \, |\widehat{\psi}(\lambda)|^2 \, d^2\lambda$$

with $\widehat{\psi}(\lambda)$ the Fourier transform of $\psi(x)$. For every $a > 0$, if $b = \frac{1}{2a}$ then $\lambda_j^2 \leqslant (a|\lambda|^2 + b)^2$. It follows that $\left\|(A_\Gamma)_j \partial_{x_j} \psi\right\|^2_{L^2} \leqslant C\|(a|\lambda|^2 + b) \, \widehat{\psi}\|^2_{L^2}$ which implies that for all $a' > 0$ (arbitrary small) there exists a b' (depending on a') such that

$$\left\|(A_\Gamma)_j \partial_{x_j} \psi\right\|^2_{L^2} \leqslant a' \, \||\lambda|^2 \, \widehat{\psi}\|^2_{L^2} + b' \, \|\widehat{\psi}\|^2_{L^2} = a' \, \|\Delta_x \psi\|^2_{L^2} + b' \, \|\psi\|^2_{L^2}.$$

This inequality implies that T_1 is infinitesimally bounded with respect to $-{}^1\!/\!{}_2\Delta_x$ and the thesis follows form the *Kato-Rellich Theorem* (see [RS75] Theorem X.12).
- *Step 2.* The Bloch-Landau Hamiltonian is

$$H_{\mathrm{BL}} := \frac{1}{2}\left[-i\nabla_x - A_\Gamma(x) - A(x)\right]^2 + V_\Gamma(x) + \phi(x). \tag{123}$$

Assumptions ($\mathrm{A_w}$) and (B) imply that $(A_\Gamma + A)_j \in C^1(\mathbb{R}^2, \mathbb{R})$, $j = 1, 2$, and $V_\Gamma + \phi \in L^2_{\mathrm{loc}}(\mathbb{R}^2)$ and this assures that H_{BL} is essentially self-adjoint on $C^\infty_{\mathrm{c}}(\mathbb{R}^2, \mathbb{C})$ (see [RS75] Theorem X.34). Let $A = A_0 + A_B$ be the decomposition of the external vector potential with A_0 smooth and bounded and $A_B = \frac{1}{2}(-x_2, x_1)$. By posing $D := \nabla_x - iA_B$, $\Delta_M := |D|^2$ and $A_{\mathrm{b}} := A_\Gamma + A_0$, the Hamiltonian H_{BL} reads

$$H_{\mathrm{BL}} = -\frac{1}{2}\Delta_M - A_{\mathrm{b}} \cdot D + \frac{1}{2}T.$$

where $T := i(\nabla_x \cdot A_{\mathrm{b}}) + |A_{\mathrm{b}}|^2 + 2(V_\Gamma + \phi)$. The operator $T - 2V_\Gamma$ is bounded and the observation that $A_{\mathrm{b}} \cdot D$ is infinitesimally bounded with respect to $-{}^1\!/\!{}_2\Delta_M$ is an immediate consequence of Lemma A.2. the assumption $\int_{M_\Gamma} |V_\Gamma(x)|^2 \, d^2x < +\infty$ implies that V_Γ is *uniformly locally* L^2 and hence, infinitesimally bounded with respect to $-\Delta_x$ (see Theorem XIII.96 in [RS78]). As proved in [AHS78] (Theorem 2.4) this is enough to claim that V_Γ is aslo infinitesimally bounded with respect to $-\Delta_M$. Therefore, by the *Kato-Rellich Theorem* it follows that the domain of self-adjointness of H_{BL} coincides with the domain of self-adjointness of the magnetic-Laplacian, which is $\mathcal{H}^2_{\mathrm{M}}(\mathbb{R}^2)$. ■

A.2 Band spectrum of H_{per}

We describe the spectral properties of the periodic Hamiltonian. The Bloch-Floquet transform maps unitarily H_{per} in $H^{\mathcal{Z}}_{\mathrm{per}} := \int^{\oplus}_{M^*_\Gamma} H_{\mathrm{per}}(k) \, d^2\underline{k}$. Then to have information about the spectrum of H_{per} we need to study the spectra of the family of Hamiltonians

$$H_{\mathrm{per}}(k) = \frac{1}{2}\left[-i\nabla_\theta + k - A_\Gamma(\theta)\right]^2 + V_\Gamma(\theta) = -\frac{1}{2}\Delta_\theta + T_1(k) + \frac{1}{2}T_2(k)$$

where $T_1(k) := i(A_\Gamma - \frac{k}{2}) \cdot \nabla_\theta$ and $T_2(k) := i(\nabla_\theta \cdot A_\Gamma) + |k|^2 + |A_\Gamma|^2 + 2V_\Gamma$ are operators acting on the Hilbert space $\mathcal{H}_\mathrm{f} := L^2(\mathbb{V}, d^2\theta)$ with $\mathbb{V} := \mathbb{R}^2/\Gamma$ (*Voronoi torus*).

Proof of Proposition 3.3.
- (i) The operator $-{}^1\!/_2\Delta_\theta$ on the Hilbert space $\mathcal{H}_\mathrm{f}$, is essentially selfadjoint on $C^\infty(\mathbb{V})$, has domain of self-adjointness $\mathcal{D} := \mathcal{H}^2(\mathbb{V})$ and its spectrum is pure point with $\{\mathrm{e}^{i\theta\cdot\gamma^*}\}_{\gamma^*\in\Gamma^*}$ a complete orthogonal system of eigenvectors. If Assumption ($\mathrm{A_w}$) holds true then $T_2(k)$ infinitesimally bounded with respect to $-{}^1\!/_2\Delta_\theta$, indeed $T - 2V_\Gamma$ is bounded and V_Γ is infinitesimally bounded (see [RS78] Theorem XIII.97). With a Fourier estimate similar to those in the proof of Proposition 2.1 one can also show that $T_1(k)$ is infinitesimally bounded with respect to $-{}^1\!/_2\Delta_\theta$, hence the *Kato-Rellich Theorem* implies that $H_\mathrm{per}(k)$ is essentially self-adjoint on $C^\infty(\mathbb{T}^2)$ and self-adjoint on the domain $\mathcal{D}$. Moreover since $-{}^1\!/_2\Delta_\theta$ is bounded below then also $H_\mathrm{per}(k)$ is bounded below.
- (ii) For all ζ in the resolvent set of $-{}^1\!/_2\Delta_\theta$ the *resolvent operator* $r_0(\zeta) := (-{}^1\!/_2\Delta_\theta - \zeta\mathbb{1}_{\mathcal{H}_\mathrm{f}})^{-1}$ is a compact operator. Since $T_1(k) + \frac{1}{2}T_2(k)$ is a bounded perturbation of $-{}^1\!/_2\Delta_\theta$ it follows that $H_\mathrm{per}(k)$ has compact resolvent (see [RS78] Theorem XIII.68) and moreover it has a purely discrete spectrum with eigenvalues $\mathcal{E}_n(k) \to +\infty$ as $n \to +\infty$ (see [RS78] Theorem XIII.64).
- (iii) The continuity of the function $\mathcal{E}_n(\cdot)$ follows from the *perturbation theory of discrete spectrum* (see [RS78] Theorem XII.13). Indeed, as discussed in Remark 3.2, $H_\mathrm{per}(\cdot)$ is an *analytic family (of type A) in the sense of Kato*. Finally, since $H_\mathrm{per}(k-\gamma^*) = \tau(\gamma^*)H_\mathrm{per}(k)\tau(\gamma^*)^{-1}$, with $\tau(\gamma^*)$ a unitary operator, then $\mathcal{E}_n(\cdot)$ are Γ^*-periodic. ■

A.3 The Landau Hamiltonian H_L

The *Landau Hamiltonian* is the operator

$$H_\mathrm{L} := -\frac{1}{2}\Delta_M = \frac{1}{2}\left(K_1^2 + K_2^2\right) = \frac{1}{2}\left[\left(-i\frac{\partial}{\partial x_1} + \frac{1}{2}x_2\right)^2 + \left(-i\frac{\partial}{\partial x_2} - \frac{1}{2}x_1\right)^2\right] \tag{124}$$

where $K_j := -iD_j$, with $j = 1, 2$, are the *kinetic momenta.* The Landau Hamiltonian H_L is essentially self-adjoint on on $C^\infty_\mathrm{c}(\mathbb{R}^2;\mathbb{C}) \subset L^2(\mathbb{R}^2)$ (see [RS75] Theorem X.34) and its domain of self-adjointness is exactly the second magnetic-Sobolev space $\mathcal{H}^2_\mathrm{M}(\mathbb{R}^2)$ defined in Section A.1. To describe the spectrum of H_L is helpful to introduce another pair of operators: $G_1 := -i\partial_{x_1} - \frac{1}{2}x_2$ and $G_2 := i\partial_{x_2} - \frac{1}{2}x_1$. The operators K_1, K_2, G_1, G_2 are all essentially self-adjoint on $C^\infty_\mathrm{c}(\mathbb{R}^2;\mathbb{C})$ (they have *deficiency indices* equal to zero) and on this domain the following commutation relations hold true

$$[K_1; K_2] = [G_1; G_2] = i\mathbb{1}, \quad [G_j; K_i] = 0. \tag{125}$$

The last of the (124) implies $[G_j; H_\mathrm{L}] = 0$ hence the operators G_1 and G_2 are cause for the degeneration of the spectral eigenspaces of H_L. It is a common lore to introduce the *annihilation operator* $\mathfrak{a} := {}^i\!/\!\sqrt{2}(K_2 - iK_1)$ (its adjoint $\mathfrak{a}^\dagger$ is called *creation operator*) and the *degeneration operator* $\mathfrak{g} := {}^i\!/\!\sqrt{2}(G_2 - iG_1)$. They fulfill the following (bosonic) commutation relation

$$[\mathfrak{a}; \mathfrak{a}^\dagger] = [\mathfrak{g}; \mathfrak{g}^\dagger] = \mathbb{1}, \quad [\mathfrak{g}; H_\mathrm{L}] = [\mathfrak{g}^\dagger; H_\mathrm{L}] = 0, \quad [\mathfrak{a}; H_\mathrm{L}] = \mathfrak{a}, \quad [\mathfrak{a}^\dagger; H_\mathrm{L}] = -\mathfrak{a}^\dagger. \tag{126}$$

The last two relations follow from the equality $H_{\mathrm{L}} = \mathfrak{a}\mathfrak{a}^{\dagger} - {}^{1}\!/_{2}\mathbb{1} = \mathfrak{a}^{\dagger}\mathfrak{a} + {}^{1}\!/_{2}\mathbb{1}$. Define the *ground state* $\psi_0 \in L^2(\mathbb{R}^2)$ as the normalized solution of $\mathfrak{g}\psi_0 = 0 = \mathfrak{a}\psi_0$, i.e. $\psi_0(x) = C\mathrm{e}^{-\frac{1}{4}|x|^2}$. The *generalized Hermite function* of order (n, m) is defined to be $\psi_{n,m} := \frac{1}{\sqrt{n!m!}} \left(\mathfrak{g}^{\dagger}\right)^m \left(\mathfrak{a}^{\dagger}\right)^n \psi_0$. We will denote by $\mathcal{F} \subset L^2(\mathbb{R}^2)$ the set of the finite linear combinations of the vectors $\psi_{n,m}$ and we will call it the *Hermite domain*. Clearly $\mathcal{F} \subset S(\mathbb{R}^2)$ (the Schwartz space).

Lemma A.1. *With the notation above:*

(i) *the set* $\{\psi_{n,m} \ : \ n, m = 0, 1, 2, \ldots\}$ *is a complete orthonormal basis for* $L^2(\mathbb{R}^2)$ *and so* $\mathcal{F}$ *is a dense domain;*

(ii) *the spectrum of* H_L *is pure point and is given by* $\{\lambda_n := (n + {}^{1}\!/_{2}) \ : \ n = 0, 1, 2, \ldots\}$, *moreover* $H_L\ \psi_{n,m} = \lambda_n\ \psi_{n,m}$ *for every* $m = 0, 1, 2, \ldots$ *(degeneration index);*

(iii) H_L *is essentially self-adjoint on* $\mathcal{F}$ *and the closure of* $\mathcal{F}$ *with respect to the magnetic-Sobolev norm coincides with the magnetic-Sobolev space* $\mathcal{H}^2_{\mathrm{M}}(\mathbb{R}^2)$.

Proof. - (i) Let $\mathcal{W} : L^2(\mathbb{R}^2, d^2x) \to L^2(\mathbb{R}, du) \otimes L^2(\mathbb{R}, dv)$ be the unitary map such that the conjugate pairs (K_1, K_2) and (G_1, G_2) are transformed by $\mathcal{W} \ldots \mathcal{W}^{-1}$ into the canonical pairs $(u, -i\partial_u)$ and $(v, -i\partial_v)$. The existence of such a unitary $\mathcal{W}$ will be discussed in Appendix B. Obviously $\mathfrak{a} \mapsto \widetilde{\mathfrak{a}} = {}^{1}\!/_{\sqrt{2}}(u + \partial_u)$, $\mathfrak{g} \mapsto \widetilde{\mathfrak{g}} = {}^{1}\!/_{\sqrt{2}}(v + \partial_v)$ and $\widetilde{\psi}_0 := \mathcal{W}\psi_0$ is the solution of $\widetilde{\mathfrak{g}}\widetilde{\psi}_0 = \widetilde{\mathfrak{a}}\widetilde{\psi}_0 = 0$, namely $\widetilde{\psi}_0(u, v) = h_0(u) \otimes h_0(v)$ where $h_0(t) := \pi^{-\frac{1}{4}}\mathrm{e}^{-\frac{1}{2}t^2}$ is the 0-th Hermite function. Then $\widetilde{\psi}_{n,m}(u, v) := (\mathcal{W}\psi_{n,m})(u, v) = h_n(u) \otimes h_m(v)$ which shows that the functions $\widetilde{\psi}_{n,m}$ define an orthonormal basis for $L^2(\mathbb{R}, du) \otimes L^2(\mathbb{R}, dv)$ since the Hermite functions h_n are an orthonormal system for $L^2(\mathbb{R})$. The claim follows since $\mathcal{W}$ is a unitary map.
- (ii) Clearly $H_{\mathrm{L}}\psi_0 = (\mathfrak{a}^{\dagger}\mathfrak{a} + {}^{1}\!/_{2}\mathbb{1})\psi_0 = {}^{1}\!/_{2}\psi_0$ and from relations (126) it follows that $H_{\mathrm{L}}\psi_{n,m} = \frac{1}{\sqrt{n!m!}} \left(\mathfrak{g}^{\dagger}\right)^m H_{\mathrm{L}} \left(\mathfrak{a}^{\dagger}\right)^n \psi_0 = {}^{1}\!/_{2}\psi_{n,m} + \frac{1}{\sqrt{n!m!}} \left(\mathfrak{g}^{\dagger}\right)^m \left(\mathfrak{a}^{\dagger}\mathfrak{a}\right) \left(\mathfrak{a}^{\dagger}\right)^n \psi_0 = \lambda_n\ \psi_{n,m}$. Then the generalized Hermite functions $\psi_{n,m}$ are a complete set of orthonormal eigenvectors for H_{L}. This proves that the spectrum of H_{L} is pure point.
- (iii) The operator H_{L} is essentially self-adjoint in $\mathcal{F}$ since the deficiency indices are both zero. This implies that last part of the claim. ■

Lemma A.2. *The operators* K_1, K_2, $\mathfrak{a}$ *and* $\mathfrak{a}^{\dagger}$ *are infinitesimally bounded with respect to* H_L.

Proof. Since $K_1 = {}^{1}\!/_{\sqrt{2}}(\mathfrak{a} + \mathfrak{a}^{\dagger})$ and $K_2 = {}^{1}\!/_{i\sqrt{2}}(\mathfrak{a} - \mathfrak{a}^{\dagger})$ it is enough to prove the claim for $\mathfrak{a}$ and $\mathfrak{a}^{\dagger}$. Let $\psi := \sum_{n,m=0}^{+\infty} c_{n,m}\ \psi_{n,m} \in \mathcal{H}^2_{\mathrm{M}}(\mathbb{R}^2)$. An easy computation shows that

$$\|\mathfrak{a}\psi\|^2_{L^2} = \sum_{n,m=0}^{+\infty} |c_{n,m}|^2\ n, \qquad\qquad \|a^{\dagger}\psi\|^2_{L^2} = \sum_{n,m=0}^{+\infty} |c_{n,m}|^2\ (n+1)\,.$$

Since $n \leqslant n + 1 \leqslant 2\left(n + \frac{1}{2}\right) \leqslant a\left(n + \frac{1}{2}\right)^2 + \frac{1}{a}$ holds true for any $a > 0$ (arbitrarily small), then

$$\|\mathfrak{a}^{\sharp}\psi\|^2_{L^2} \leqslant a \sum_{n,m=0}^{+\infty} |c_{n,m}|^2 \left(n + \frac{1}{2}\right)^2 + b \sum_{n,m=0}^{+\infty} |c_{n,m}|^2 = a\|H_{\mathrm{L}}\psi\|^2_{L^2} + b\|\psi\|^2_{L^2}$$

with $b := \frac{1}{a} + \frac{1}{2}$ where $\mathfrak{a}^\sharp$ denotes either $\mathfrak{a}$ or $\mathfrak{a}^\dagger$. ■

B Canonical transform for fast and slow variables

This appendix is devoted to the concrete realization of the *von Neumann unitary* $\mathcal{W}$ introduced (in abstract way) in Section 4.2. The unitary $\mathcal{W}$ maps the fast and slow variables, which satisfy canonical commutation relation, into a set of canonical Schrödinger operators. In Section B.1 we derive a general version of the transform $\mathcal{W}$ "by hand", as a composition of three sequential transforms. In Section B.2 we compute the integral kernel of $\mathcal{W}$.

B.1 The transform $\mathcal{W}$ built "by hand"

Let $\mathcal{H} := L^2(\mathbb{R}^2, d^2r)$ be the initial Hilbert space, with $r := (r_1, r_2)$. Let $Q_r := (Q_{r_1}, Q_{r_2})$ where Q_{r_j} is the multiplication operator by r_j and $P_r := (P_{r_1}, P_{r_2})$ where $P_{r_1} := -i\hbar\partial_{r_j}$, with $j = 1, 2$. Consider the *fast* and *slow* operators

$$\text{(fast)} \begin{cases} K_1 := -\dfrac{\alpha}{2\beta}\, v \cdot Q_r - \dfrac{\alpha\beta}{\hbar}\, w^* \cdot P_r \\ K_2 := \dfrac{\alpha}{2\beta}\, w \cdot Q_r - \dfrac{\alpha\beta}{\hbar}\, v^* \cdot P_r \end{cases} \qquad \text{(slow)} \begin{cases} G_1 := \dfrac{1}{2}\, v \cdot Q_r - \dfrac{\beta^2}{\hbar}\, w^* \cdot P_r \\ G_2 := \dfrac{1}{2}\, w \cdot Q_r + \dfrac{\beta^2}{\hbar}\, v^* \cdot P_r \end{cases} \tag{127}$$

with $\alpha, \beta \in \mathbb{C}$ and $v, w, v^*, w^* \in \mathbb{R}^2$ such that $v \cdot v^* = w \cdot w^* = 1$, $v^* \cdot w = v \cdot w^* = 0$ and $|v \wedge w| = \ell^2 > 0$.

Remark B.1. The choice $v = b^*$, $w = a^*$, $\alpha = \sqrt{\iota_q}$ and $\beta = \sqrt{\iota_q}\,\delta$ defines the operators (56), while the choice $v = v^* = (0, -1)$, $w = w^* = (-1, 0)$, $\alpha = \beta = 1$ defines the *kinetic momenta* and the related conjugate operators introduced in Section A.3. ♦◊

Observing that $[a \cdot Q_r + b \cdot P_r ; c \cdot Q_r + d \cdot P_r] = i\hbar(a \cdot d - b \cdot c)\mathbb{1}_{\mathcal{H}}$ one deduce that the operators (127) verify the following *canonical commutation relations* (CCR)

$$[K_1, K_2] = i\alpha^2 \mathbb{1}_{\mathcal{H}}, \qquad [G_1, G_2] = i\beta^2 \mathbb{1}_{\mathcal{H}}, \qquad [K_i, G_j] = 0, \quad i, j = 1, 2. \tag{128}$$

The *Stone-von Neumann uniqueness theorem* (see [BR97] Corollary 5.2.15) assures the existence of a unitary map $\mathcal{W}$ (*von Neumann unitary*)

$$\mathcal{W} : \mathcal{H} \longrightarrow \mathcal{H}_{\mathrm{w}} := \mathcal{H}_{\mathrm{s}} \otimes \mathcal{H}_{\mathrm{f}} := L^2(\mathbb{R}, dx_{\mathrm{s}}) \otimes L^2(\mathbb{R}, dx_{\mathrm{f}}) \tag{129}$$

such that

$$\mathcal{W} G_1 \mathcal{W}^{-1} := Q_{\mathrm{s}} = \text{multiplication by } x_{\mathrm{s}}, \qquad \mathcal{W} G_2 \mathcal{W}^{-1} := P_{\mathrm{s}} = -i\beta^2 \frac{\partial}{\partial x_{\mathrm{s}}} \tag{130}$$

$$\mathcal{W} K_1 \mathcal{W}^{-1} := Q_{\mathrm{f}} = \text{multiplication by } x_{\mathrm{f}}, \qquad \mathcal{W} K_2 \mathcal{W}^{-1} := P_{\mathrm{f}} = -i\alpha^2 \frac{\partial}{\partial x_{\mathrm{f}}}. \tag{131}$$

In other words, $(Q_{\mathrm{s}}, P_{\mathrm{s}})$ is a pair of operators which defines a Schrödinger representation on the Hilbert space $\mathcal{H}_{\mathrm{s}} := L^2(\mathbb{R}, dx_{\mathrm{s}})$ while the pair $(Q_{\mathrm{f}}, P_{\mathrm{f}})$ defines a Schrödinger representation on the Hilbert space $\mathcal{H}_{\mathrm{f}} := L^2(\mathbb{R}, dx_{\mathrm{f}})$. Our purpose is to give an explicit construction for $\mathcal{W}$. First of all consider the change of coordinates $(r_1, r_2) \mapsto (k_1 := \frac{v \cdot r}{\ell}, k_2 := \frac{w \cdot r}{\ell})$. The inverse transforms are defined by $r_1(k) = \frac{1}{\ell}(w_2 k_1 - v_2 k_2)$ and $r_2(k) = \frac{1}{\ell}(v_1 k_2 - w_1 k_1)$. The map $\mathcal{J} : L^2(\mathbb{R}^2, d^2 r) \to L^2(\mathbb{R}^2, d^2 k)$ defined by $(\mathcal{J}\psi)(k) := \psi(r(k))$ if $\psi \in L^2(\mathbb{R}^2, dr)$ is unitary since the change of coordinates is invertible and isometric. Moreover $\mathcal{J} Q_{r_j} \mathcal{J}^{-1}$ acts on $L^2(\mathbb{R}^2, dk)$ as the multiplication by $r_j(k)$, while $\mathcal{J} P_{r_j} \mathcal{J}^{-1} = \frac{v_j}{\ell} P_{k_1} + \frac{w_j}{\ell} P_{k_2}$ where $P_{k_j} := -i\hbar \partial_{k_j}$, with $j = 1, 2$. Then

$$\mathcal{J} G_1 \mathcal{J}^{-1} := \quad \frac{\ell}{2}\, Q_{k_1} - \frac{\beta^2}{\hbar}\frac{1}{\ell}\, P_{k_2}, \qquad \mathcal{J} G_2 \mathcal{J}^{-1} := \quad \frac{\ell}{2}\, Q_{k_2} + \frac{\beta^2}{\hbar}\frac{1}{\ell}\, P_{k_1} \tag{132}$$

$$\mathcal{J} K_1 \mathcal{J}^{-1} := -\frac{\alpha}{\beta}\frac{\ell}{2}\, Q_{k_1} - \frac{\alpha\beta}{\hbar}\frac{1}{\ell}\, P_{k_2}, \qquad \mathcal{J} K_2 \mathcal{J}^{-1} := \frac{\alpha}{\beta}\frac{\ell}{2}\, Q_{k_2} - \frac{\alpha\beta}{\hbar}\frac{1}{\ell}\, P_{k_1}. \tag{133}$$

Let $\mathcal{F}_{2,\mu} : L^2(\mathbb{R}, dk_2) \to L^2(\mathbb{R}, d\zeta_2)$ be the k_2-*Fourier transfor* of weight μ, defined by $(\mathcal{F}_{2,\mu}\psi)(\zeta_2) := \sqrt{\frac{|\mu|}{2\pi}} \int_{\mathbb{R}} \mathrm{e}^{-i\mu\zeta_2 k_2} \psi(k_2)\, dk_2$ and let $\Pi_1 : L^2(\mathbb{R}, dk_1) \to L^2(\mathbb{R}, d\zeta_1)$ be the k_1-*parity operator* defined by $(\Pi_2\psi)(\zeta_1) := \psi(-\zeta_1)$ (namely by the change of coordinates $k_1 \mapsto \zeta_1$). Let $\mathbb{I}$ the unitary map which identifies $L^2(\mathbb{R}^2, d^2k)$ with $L^2(\mathbb{R}, dk_1) \otimes L^2(\mathbb{R}, dk_2)$. Let $Q_\zeta := (Q_{\zeta_1}, Q_{\zeta_2})$ whit Q_{ζ_j} the multiplication operator by ζ_j and $P_\zeta := (P_{\zeta_1}, P_{\zeta_2})$ where $P_{\zeta_1} := -i\hbar\partial_{\zeta_j}$, with $j = 1, 2$. One can check that

$$\mathcal{F}_{2,\mu} Q_{k_2} \mathcal{F}_{2,\mu}^{-1} = -\frac{1}{\mu\hbar} P_{\zeta_2}, \quad \mathcal{F}_{2,\mu} P_{k_2} \mathcal{F}_{2,\mu}^{-1} = \mu\hbar Q_{\zeta_2}, \quad \Pi_1 Q_{k_1} \Pi_1^{-1} = -Q_{\zeta_1}, \quad \Pi_1 P_{k_1} \Pi_1^{-1} = -P_{\zeta_1}. \tag{134}$$

Fix $\mu := -\frac{\ell^2}{2\beta^2}$, then the unitary map $\mathcal{L} := (\Pi_1 \otimes \mathcal{F}_{2,\mu}) \circ \mathbb{I} \circ \mathcal{J} : \mathcal{H} \to L^2(\mathbb{R}, d\zeta_1) \otimes L^2(\mathbb{R}, d\zeta_2)$ acts on the operators (127) in the following way

$$\mathcal{L} G_1 \mathcal{L}^{-1} := -\frac{\ell}{2}\left(Q_{\zeta_1} - Q_{\zeta_2}\right), \qquad \mathcal{L} G_2 \mathcal{J}^{-1} := -\frac{1}{\ell}\frac{\beta^2}{\hbar}\left(P_{\zeta_1} - P_{\zeta_2}\right) \tag{135}$$

$$\mathcal{L} K_1 \mathcal{L}^{-1} := \frac{\alpha}{\beta}\frac{\ell}{2}\left(Q_{\zeta_1} + Q_{\zeta_2}\right), \qquad \mathcal{L} K_2 \mathcal{L}^{-1} := \quad \frac{1}{\ell}\frac{\alpha\beta}{\hbar}\left(P_{\zeta_1} + P_{\zeta_2}\right). \tag{136}$$

Now we can consider the change of coordinates $(\zeta_1, \zeta_2) \mapsto (x_{\mathrm{s}}, x_{\mathrm{f}})$ defined by

$$\begin{cases} x_{\mathrm{s}} = -\dfrac{\ell}{2}(\zeta_1 - \zeta_2) \\ x_{\mathrm{f}} = \quad \dfrac{\alpha}{\beta}\dfrac{\ell}{2}(\zeta_1 + \zeta_2) \end{cases} \qquad \begin{cases} \zeta_1 = -\dfrac{1}{\ell}\left(x_{\mathrm{s}} - \dfrac{\beta}{\alpha}\, x_{\mathrm{f}}\right) \\ \zeta_2 = \quad \dfrac{1}{\ell}\left(x_{\mathrm{s}} + \dfrac{\beta}{\alpha}\, x_{\mathrm{f}}\right) \end{cases}$$

The jacobian of this transformation is $|\partial(\zeta_1, \zeta_2)/\partial(x_{\mathrm{s}}, x_{\mathrm{f}})| = \frac{2}{\ell^2}\left|\frac{\beta}{\alpha}\right| =: C$, then the map $(\mathcal{R}\psi)(x_{\mathrm{s}}, x_{\mathrm{f}}) := \sqrt{C}\ \psi(\zeta(x_{\mathrm{s}}, x_{\mathrm{f}}))$ defines a unitary map $\mathcal{R} : L^2(\mathbb{R}^2, d^2\zeta) \to L^2(\mathbb{R}^2, dx_{\mathrm{s}}\ dx_{\mathrm{f}})$. With a direct computation one can check that $\mathcal{R} Q_{\zeta_j} \mathcal{R}^{-1}$ acts on $L^2(\mathbb{R}^2, dx_{\mathrm{s}}\ dx_{\mathrm{f}})$ as the multiplication by $\zeta_j(x_{\mathrm{s}}, x_{\mathrm{f}})$, while $\mathcal{R} P_{\zeta_j} \mathcal{R}^{-1} = \frac{\ell\hbar}{2\alpha\beta}\left(\frac{(-1)^j\alpha}{\beta} P_{\mathrm{s}} + P_{\mathrm{f}}\right)$, with $j = 1, 2$. This shows that the unitary map $\mathcal{W} := \mathbb{I} \circ \mathcal{R} \circ \mathbb{I}^{-1} \circ \mathcal{L} := \mathbb{I} \circ \mathcal{R} \circ \mathbb{I}^{-1} \circ (\Pi_1 \otimes \mathcal{F}_{2,\mu}) \circ \mathbb{I} \circ \mathcal{J}$ is the von Neumann unitary that verifies the relations (130) and (131).

B.2 The integral kernel of $\mathcal{W}$

The unitary operators $\mathcal{J} : L^2(\mathbb{R}^2, d^2k) \to L^2(\mathbb{R}^2, d^2r)$ and $\mathcal{R} : L^2(\mathbb{R}^2, d^2\zeta) \to L^2(\mathbb{R}^2, dx_{\mathrm{s}}dx_{\mathrm{f}})$ related to the change of coordinates $(r_1, r_2) \mapsto (k_1, k_2)$ and $(\zeta_1, \zeta_2) \mapsto (x_{\mathrm{s}}, x_{\mathrm{f}})$ can be written an integral operators

$$(\mathcal{J}\psi)(k) = \int_{\mathbb{R}^2} J(r; k)\ \psi(r)\ d^2r, \qquad (\mathcal{R}\varphi)(x_{\mathrm{s}}, x_{\mathrm{f}}) = \int_{\mathbb{R}^2} R(\zeta; x_{\mathrm{s}}, x_{\mathrm{f}})\ \varphi(\zeta)\ d^2\zeta$$

with distributional integral kernels

$$J(r_1, r_2; k_1, k_2) := \delta\left(r_1 - \frac{1}{\ell}(w_2k_1 - v_2k_2)\right)\ \delta\left(r_2 + \frac{1}{\ell}(w_1k_1 - v_1k_2)\right)$$

$$R(\zeta_1, \zeta_2; x_{\mathrm{s}}, x_{\mathrm{f}}) := \sqrt{C}\ \delta\left(\zeta_1 + \frac{1}{\ell}\left(x_{\mathrm{s}} - \frac{\beta}{\alpha}\ x_{\mathrm{f}}\right)\right)\ \delta\left(\zeta_2 - \frac{1}{\ell}\left(x_{\mathrm{s}} + \frac{\beta}{\alpha}\ x_{\mathrm{f}}\right)\right)$$

with $C = \frac{2}{\ell^2}\left|\frac{\beta}{\alpha}\right|$. The k_1-parity operator Π_1 can be written as an integral operator with the distributional kernel $\delta(r_1 + k_1)$ while the integral kernel of the k_2-Fourier transform $\mathcal{F}_{2,\mu}$, with $\mu := -\frac{\ell^2}{2\beta^2}$, is $\frac{\ell}{2|\beta|\sqrt{\pi}}\mathrm{e}^{i\frac{\ell^2}{2\beta^2}\zeta_2k_2}$. Then the unitary map $\mathbb{I}^{-1} \circ (\Pi_1 \otimes \mathcal{F}_{2,\mu}) \circ \mathbb{I} : L^2(\mathbb{R}^2, d^2k) \to L^2(\mathbb{R}^2, d^2\zeta)$ as the integral distributional kernel

$$L(k_1, k_2; \zeta_1, \zeta_2) := \delta(\zeta_1 + k_1)\ \frac{\ell}{2|\beta|\sqrt{\pi}}\mathrm{e}^{i\frac{\ell^2}{2\beta^2}\zeta_2k_2}.$$

Summarizing the total transform $\mathcal{W} : L^2(\mathbb{R}^2, d^2r) \to L^2(\mathbb{R}^2, dx_{\mathrm{s}}\ dx_{\mathrm{f}})$ (up to the obvious identification $\mathbb{I}$) can be expressed as an integral operator

$$(\mathcal{W}\psi)(x_{\mathrm{s}}, x_{\mathrm{f}}) = \int_{\mathbb{R}^2} W(r; x_{\mathrm{s}}, x_{\mathrm{f}})\ \Psi(r)\ d^2r$$

with a (total) integral distributional kernel

$$W(r_1, r_2; x_{\mathrm{s}}, x_{\mathrm{f}}) := \frac{\ell}{\sqrt{2\pi|\alpha\beta|}}\ \delta\left(v \cdot r - \left(x_{\mathrm{s}} - \frac{\beta}{\alpha}\ x_{\mathrm{f}}\right)\right)\ \mathrm{e}^{i\frac{w\cdot r}{2\beta^2}\left(x_{\mathrm{s}} + \frac{\beta}{\alpha}\ x_{\mathrm{f}}\right)}.$$

References

[AHS78] J. E. Avron, I. Herbst, and B. Simon. Schrödinger Operators with Magnetic Fields I. General Interactions. *Duke Math. J.*, **45** (4): 847–883, 1978.

[AM76] N. W. Ashcroft and N. D. Mermin. *Solid State Physics.* Saunders, 1976.

[ASS83] J. E. Avron, R. Seiler, and B. Simon. Homotopy and Quantization in Condensed Matter Physics. *Phys. Rev. Lett.*, **51**: 51–53, 1983.

[Avr04] J. E. Avron. *Colored Hofstadter butterflies.* Ph. Blanchard, G. Dell'Antonio (eds.), Multiscale Methods in Quantum Mechanics: Theory and Experiments. Birkhäuser, 2004.

[Bel88] J. Bellissard. *C^*-algebras in solid state physics. 2D electrons in a uniform magnetic field*, volume II of *Operator algebras and applications.* University Press, 1988.

[BR97] O. Bratteli and D. W. Robinson. *Equilibrium States, Models in Quantum Statistical Mechanics*, volume II of *Operator Algebras and Quantum Statistical Mechanics.* Springer-Verlag, 1997.

[BSE94] J. Bellissard, H. Schulz-Baldes, and A. van Elst. The Non Commutative Geometry of the Quantum Hall Effect. *J. Math. Phys.*, **35**: 5373–5471, 1994.

[CTVR06] D. Ceresoli, T. Thonhauser, D. Vanderbilt, and R. Resta. Orbital magnetization in crystalline solids: multi-band insulators, Chern insulators, and metals. *Phys. Rev. B*, **74**: 024408, 2006.

[DL10] G. De Nittis and M. Lein. Applications of magnetic ΨDO techniques to Space-adiabatic Perturbation Theory. Preprint *arXive.org/1006.3103*, 2010.

[GA03] O. Gat and J. E. Avron. Semiclassical analysis and the magnetization of the Hofstadter model. *Phys. Rev. Lett.*, **91**: 186801, 2003.

[GMS91] C. Gérard, A. Martinez, and J. Sjöstrand. A mathematical approach to the effective Hamiltonian in perturbed periodic problems. *Comm. Math. Phys.*, **142**: 217–244, 1991.

[GPPO00] V. A. Geyler, I. Yu. Popov, A. V. Popov, and A. A. Ovechkina. Fractal spectrum of periodic quantum systems in a magnetic field. *Chaos, solitons and fractals*, **11**: 281–288, 2000.

[Gra07] G. M. Graf. Aspects of the Integer Quantum Hall Effect. *Proceedings of Symposia in Pure Mathematics*, **76.1**: 429–442, 2007.

[Har55] P. G. Harper. Single band motion of conduction electrons in a uniform magnetic field. *Proc. Phys. Soc. A*, **68**: 874–892, 1955.

[Hof76] D. R. Hofstadter. Energy levels and wave functions of Bloch electrons in rational and irrational magnetic fields. *Phys. Rev. B*, **14**: 2239–2249, 1976.

[HS88] B. Helffer and J. Sjöstrand. Analyse semi-classique pour l'équation de Harper (avec application à l'étude de Schrödinger avec champ magnétique). *Mém. Soc. Math. France*, **34**: 1761–1771, 1988.

[HS89a] B. Helffer and J. Sjöstrand. Analyse semi-classique pour l'équation de Harper III. *Mém. Soc. Math. France*, **39**, 1989.

[HS89b] B. Helffer and J. Sjöstrand. *Équation de Schrödinger avec champ magnétique et équation de Harper*, volume **345** of *Lecture Notes in Physics.* Springer-Verlag, 1989.

[HS90] B. Helffer and J. Sjöstrand. Analyse semi-classique pour l'équation de Harper II (comportement semi-classique près d'un rationnel). *Mém. Soc. Math. France*, **40**, 1990.

[IMP07] V. Iftimie, M. Măntoiu, and R. Purice. Magnetic pseudodifferential operators. *Publ. RIMS.*, **43** no. 3: 585–623, 2007.

[IMP09] V. Iftimie, M. Măntoiu, and R. Purice. The magnetic formalism; new results. *Contemporary Mathematics*, **500**: 123–138, 2009.

[Kuc93] P. Kuchment. *Floquet Theory for Partial Differential Equations.* Operator Theory: Advances and Applications. Birkhäuser, 1993.

[Las94] Y. Last. Zero measure spectrum for the almost Mathieu operator. *Comm. Math. Phys.*, **164**: 421–432, 1994.

[Las05] Y. Last. *Spectral theory of Sturm-Liouville operators on infinite intervals: A review of recent developments.* Sturm-Liouville Theory (W. O. Amrein, A. M. Hinz, and D. B. Pearson, eds.). Birkhäuser, 2005.

[LL01] E. H. Lieb and M. Loss. *Analysis.* American Mathematical Society, 2001.

[Mor88] G. Morandi. *Quantum Hall Effect.* Bibliopolis, 1988.

[MP04] M. Măntoiu and R. Purice. The Magnetic Weyl calculus. *J. Math. Phys.*, **45**: 1394, 2004.

[MPR05] M. Măntoiu, R. Purice, and S. Richard. Twisted Crossed Products and Magnetic Pseudodifferential Operators. *in Advances in Operator Algebras and Mathematical Physics*, pages 137–172, 2005.

[Nen02] G. Nenciu. On asymptotic perturbation theory for Quantum Mechanics: Almost invariant subspaces and gauge invariant magnetic perturbation theory. *J. Math. Phys.*, **43**: 1273–1298, 2002.

[NS04] G. Nenciu and V. Sordoni. Semiclassic limit for multistate Klein-Gordon systems: almost invariant subspaces, and scattering theory. *J. Math. Phys.*, **45**: 3676, 2004.

[OA01] D. Osadchy and J. E. Avron. Hofstadter butterfly as quantum phase diagram. *J. Math. Phys.*, **42**: 5665–5671, 2001.

[OGM08] P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin. Theory of anomalous quantum Hall effects in graphene. *Phys. Rev. B*, **77**: 195430, 2008.

[Pan07] G. Panati. Triviality of Bloch and Bloch-Dirac bundles. *Ann. Henri Poincaré*, **8**: 995–1011, 2007.

[Pei33] R. Peierls. Zur Theorie des Diamagnetismus von Leitungselektronen. *Z. Phys.*, **80**: 763–791, 1933.

[PST03a] G. Panati, H. Spohn, and S. Teufel. Effective dynamics for Bloch electrons: Peierls substitution and beyond. *Comm. Math. Phys.*, **242**: 547–578, 2003.

[PST03b] G. Panati, H. Spohn, and S. Teufel. Space-Adiabatic Perturbation Theory. *Adv. Theor. Math. Phys.*, **7**: 145–204, 2003.

[Rau74] A. Rauh. Degeneracy of Landau levels in crystals. *Phys. Status Solidi B*, **65**: K131–K135, 1974.

[Rau75] A. Rauh. On the broadening of Landau levels in crystals. *Phys. Status Solidi B*, **69**: K9–K13, 1975.

[RS75] M. Reed and B. Simon. *Fourier Analysis, Self-Adjointness*, volume II of *Methods of Modern Mathematical Phisics*. Academic Press, 1975.

[RS78] M. Reed and B. Simon. *Analysis of Operators*, volume IV of *Methods of Modern Mathematical Phisics*. Academic Press, 1978.

[Shu94] M. A. Shubin. Discrete Magnetic Laplacian. *Commun. Math. Phys.*, **164**: 259–275, 1994.

[Sob97] A. V. Sobolev. Absolute continuity of the periodic magnetic Schrödinger operator. *Invent. Math.*, **137**: 85–112, 1997.

[TCVR05] T. Thonhauser, D. Ceresoli, D. Vanderbilt, and R. Resta. Orbital magnetization in periodic insulators. *Phys. Rev. Lett.*, **95**:137205, 2005.

[Teu03] S. Teufel. *Adiabatic Perturbation Theory in Quantum Dynamics*, volume 1821 of *Lecture Notes in Mathematics*. Springer-Verlag, 2003.

[TKNN82] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. Nijs. Quantized Hall conductance in a two-dimensional periodic potential. *Phys. Rev. Lett.*, **49**: 405–408, 1982.

[Wil87] M. Wilkinson. An exact effective Hamiltonian for a perturbed Landau level. *J. Phys. A*, **20**: 1761–1771, 1987.